

## **APPENDIX**

### **CLEAN COPY OF SUBSTITUTE SPECIFICATION**

U.S. Utility Patent Application  
Application No. 09/189,702; Filed: November 10, 1998  
For: HLA Binding Peptides and Their Uses  
Inventors: SETTE et al.  
Our Ref: 2060.0370001/EKS/M-M



## HLA BINDING PEPTIDES AND THEIR USES

### REFERENCE TO RELATED APPLICATIONS

This application is a continuation-in-part of U.S. Application No. 09/098,584, filed June 17, 1998, now abandoned.

5

### REFERENCE TO A SEQUENCE LISTING AND TABLES SUBMITTED ON A COMPACT DISC

The Substitute Sequence Listing written in file Sequence Listing.txt, 92,000 bytes, created on June 22, 2005 on compact disc for Application No. 09/189,702, 10 Sette *et al.*, HLA Binding Peptides and Their Uses, is herein incorporated-by-reference.

### BACKGROUND OF THE INVENTION

The present invention relates to compositions and methods for preventing, treating or diagnosing a number of pathological states such as viral diseases and cancers.

15 In particular, it provides novel peptides capable of binding selected major histocompatibility complex (MHC) molecules and inducing an immune response.

MHC molecules are classified as either Class I or Class II molecules.

Class II MHC molecules are expressed primarily on cells involved in initiating and sustaining immune responses, such as T lymphocytes, B lymphocytes, macrophages, etc.

20 Class II MHC molecules are recognized by helper T lymphocytes and induce proliferation of helper T lymphocytes and amplification of the immune response to the particular immunogenic peptide that is displayed. Class I MHC molecules are expressed on almost all nucleated cells and are recognized by cytotoxic T lymphocytes (CTLs), which then destroy the antigen-bearing cells. CTLs are particularly important in tumor rejection and 25 in fighting viral infections.

The CTL recognizes the antigen in the form of a peptide fragment bound to the MHC class I molecules rather than the intact foreign antigen itself. The antigen must normally be endogenously synthesized by the cell, and a portion of the protein antigen is degraded into small peptide fragments in the cytoplasm. Some of these small 30 peptides translocate into a pre-Golgi compartment and interact with class I heavy chains to facilitate proper folding and association with the subunit  $\beta 2$  microglobulin. The

peptide-MHC class I complex is then routed to the cell surface for expression and potential recognition by specific CTLs.

Investigations of the crystal structure of the human MHC class I molecule, HLA-A2.1, indicate that a peptide binding groove is created by the folding of the  $\alpha 1$  and  $\alpha 2$  domains of the class I heavy chain (Bjorkman et al., Nature 329:506 (1987)). In these investigations, however, the identity of peptides bound to the groove was not determined.

Buus et al., Science 242:1065 (1988) first described a method for acid elution of bound peptides from MHC. Subsequently, Rammensee and his coworkers (Falk et al., Nature 351:290 (1991)) have developed an approach to characterize naturally processed peptides bound to class I molecules. Other investigators have successfully achieved direct amino acid sequencing of the more abundant peptides in various HPLC fractions by conventional automated sequencing of peptides eluted from class I molecules of the B type (Jardetzky, et al., Nature 353:326 (1991)) and of the A2.1 type by mass spectrometry (Hunt, et al., Science 225:1261 (1992)). A review of the characterization of naturally processed peptides in MHC Class I has been presented by Rötzschke and Falk (Rötzschke and Falk, Immunol. Today 12:447 (1991)).

Sette et al., Proc. Natl. Acad. Sci. USA 86:3296 (1989) showed that MHC allele specific motifs could be used to predict MHC binding capacity. Schaeffer et al., Proc. Natl. Acad. Sci. USA 86:4649 (1989) showed that MHC binding was related to immunogenicity. Several authors (De Bruijn et al., Eur. J. Immunol., 21:2963-2970 (1991); Pamer et al., 991 Nature 353:852-955 (1991)) have provided preliminary evidence that class I binding motifs can be applied to the identification of potential immunogenic peptides in animal models. Class I motifs specific for a number of human alleles of a given class I isotype have yet to be described. It is desirable that the combined frequencies of these different alleles should be high enough to cover a large fraction or perhaps the majority of the human outbred population.

Despite the developments in the art, the prior art has yet to provide a useful human peptide-based vaccine or therapeutic agent based on this work. The present invention provides these and other advantages.

## SUMMARY OF THE INVENTION

The present invention provides compositions comprising immunogenic peptides having binding motifs for HLA-A2.1 molecules. The immunogenic peptides,

which bind to the appropriate MHC allele, are preferably 9 to 10 residues in length and comprise conserved residues at certain positions such as positions 2 and 9. Moreover, the peptides do not comprise negative binding residues as defined herein at other positions such as positions 1, 3, 6 and/or 7 in the case of peptides 9 amino acids in length and  
 5 positions 1, 3, 4, 5, 7, 8 and/or 9 in the case of peptides 10 amino acids in length. The present invention defines positions within a motif enabling the selection of peptides which will bind efficiently to HLA A2.1.

The motifs of the inventions include peptide of 9 amino acids which have a first conserved residue at the second position from the N-terminus selected from the  
 10 group consisting of I, V, A and T and a second conserved residue at the C-terminal position selected from the group consisting of V, L, I, A and M. Alternatively, the peptide may have a first conserved residue at the second position from the N-terminus selected from the group consisting of L, M, I, V, A and T; and a second conserved residue at the C-terminal position selected from the group consisting of A and M. If the peptide  
 15 has 10 residues it will contain a first conserved residue at the second position from the N-terminus selected from the group consisting of L, M, I, V, A, and T; and a second conserved residue at the C-terminal position selected from the group consisting of V, I, L, A and M; wherein the first and second conserved residues are separated by 7 residues.

Epitopes on a number of immunogenic target proteins can be identified  
 20 using the peptides of the invention. Examples of suitable antigens include prostate cancer specific antigen (PSA), prostate specific membrane antigen (PSM), hepatitis B core and surface antigens (HBVc, HBVs) hepatitis C antigens, Epstein-Barr virus antigens, human immunodeficiency type-1 virus (HIV1), Kaposi's sarcoma herpes virus (KSHV), human papilloma virus (HPV) antigens, Lassa virus, mycobacterium tuberculosis (MT), p53 and  
 25 murine p53 (mp53), CEA, trypanosome surface antigen (TSA), members of the tyrosinase related protein (TRP) families, and Her2/neu. The peptides are thus useful in pharmaceutical compositions for both in vivo and ex vivo therapeutic and diagnostic applications.

The present invention also provides compositions comprising  
 30 immunogenic peptides having binding motifs for MHC Class I molecules. The immunogenic peptides are typically between about 8 and about 11 residues and comprise conserved residues involved in binding proteins encoded by the appropriate MHC allele. A number of allele specific motifs have been identified.

For instance, the motif for HLA-A3.2 comprises from the N-terminus to C-terminus a first conserved residue of L, M, I, V, S, A, T and F at position 2 and a second conserved residue of K, R or Y at the C-terminal end. Other first conserved residues are C, G or D and alternatively E. Other second conserved residues are H or F.

5 The first and second conserved residues are preferably separated by 6 to 7 residues.

The motif for HLA-A1 comprises from the N-terminus to the C-terminus a first conserved residue of T, S or M, a second conserved residue of D or E, and a third conserved residue of Y. Other second conserved residues are A, S or T. The first and second conserved residues are adjacent and are preferably separated from the third conserved residue by 6 to 7 residues. A second motif consists of a first conserved residue of E or D and a second conserved residue of Y where the first and second conserved residues are separated by 5 to 6 residues.

The motif for HLA-A11 comprises from the N-terminus to the C-terminus a first conserved residue of T, V, M, L, I, S, A, G, N, C D, or F at position 2 and a C-terminal conserved residue of K, R, Y or H. The first and second conserved residues are preferably separated by 6 or 7 residues.

The motif for HLA-A24.1 comprises from the N-terminus to the C-terminus a first conserved residue of Y, F or W at position 2 and a C terminal conserved residue of F, I, W, M or L. The first and second conserved residues are preferably separated by 6 to 7 residues.

Epitopes on a number of potential target proteins can be identified in this manner. Examples of suitable antigens include prostate specific antigen (PSA), prostate specific membrane antigen (PSM), hepatitis B core and surface antigens (HBVc, HBVs) hepatitis C antigens, malignant melanoma antigen (MAGE-1) Epstein-Barr virus antigens, human immunodeficiency type-1 virus (HIV1), papilloma virus antigens, Lassa virus, mycobacterium tuberculosis (MT), p53 and murine p53 (mp53), CEA, and Her2/neu, and members of the tyrosinase related protein (TRP) families. The peptides are thus useful in pharmaceutical compositions for both in vivo and ex vivo therapeutic and diagnostic applications.

30 The present invention also provides compositions comprising immunogenic peptides having binding motifs for non-A HLA alleles. The immunogenic peptides are preferably about 9 to 10 residues in length and comprise conserved residues at certain positions such as proline at position 2 and an aromatic residue (e.g., Y, W, F) or

hydrophobic residue (e.g., L, I, V, M, or A) at the carboxy terminus. In particular, an advantage of the peptides of the invention is their ability to bind to two or more different HLA alleles.

Epitopes on a number of potential target proteins can be identified in this manner. Examples of suitable antigens include prostate specific antigen (PSA), hepatitis B core and surface antigens (HBVc, HBVs) hepatitis C antigens, malignant melanoma antigen (MAGE-1) Epstein-Barr virus antigens, human immunodeficiency type-1 virus (HIV1), papilloma virus antigens, Lassa virus, mycobacterium tuberculosis (MT), p53, CEA, and Her2/neu. The peptides are thus useful in pharmaceutical compositions for both in vivo and ex vivo therapeutic and diagnostic applications.

### Definitions

The term "peptide" is used interchangeably with "oligopeptide" in the present specification to designate a series of residues, typically L-amino acids, connected one to the other typically by peptide bonds between the alpha-amino and carbonyl groups of adjacent amino acids. The oligopeptides of the invention are less than about 15 residues in length and usually consist of between about 8 and about 11 residues, preferably 9 or 10 residues.

An "immunogenic peptide" is a peptide which comprises an allele-specific motif such that the peptide will bind an MHC molecule and induce a CTL response. Immunogenic peptides of the invention are capable of binding to an appropriate HLA-A2.1 molecule and inducing a cytotoxic T cell response against the antigen from which the immunogenic peptide is derived.

Immunogenic peptides are conveniently identified using the algorithms of the invention. The algorithms are mathematical procedures that produce a score which enables the selection of immunogenic peptides. Typically one uses the algorithmic score with a "binding threshold" to enable selection of peptides that have a high probability of binding at a certain affinity and will in turn be immunogenic. The algorithm is based upon either the effects on MHC binding of a particular amino acid at a particular position of a peptide or the effects on binding of a particular substitution in a motif containing peptide.

A "conserved residue" is an amino acid which occurs in a significantly higher frequency than would be expected by random distribution at a particular position

in a peptide. Typically a conserved residue is one where the MHC structure may provide a contact point with the immunogenic peptide. At least one to three or more, preferably two, conserved residues within a peptide of defined length defines a motif for an immunogenic peptide. These residues are typically in close contact with the peptide binding groove, with their side chains buried in specific pockets of the groove itself. Typically, an immunogenic peptide will comprise up to three conserved residues, more usually two conserved residues.

As used herein, "negative binding residues" are amino acids which if present at certain positions (for example, positions 1, 3 and/or 7 of a 9-mer) will result in a peptide being a nonbinder or poor binder and in turn fail to be immunogenic i.e. induce a CTL response.

The term "motif" refers to the pattern of residues in a peptide of defined length, usually about 8 to about 11 amino acids, which is recognized by a particular MHC allele. The peptide motifs are typically different for each human MHC allele and differ in the pattern of the highly conserved residues and negative residues.

The binding motif for an allele can be defined with increasing degrees of precision. In one case, all of the conserved residues are present in the correct positions in a peptide and there are no negative residues in positions 1,3 and/or 7.

The phrases "isolated" or "biologically pure" refer to material which is substantially or essentially free from components which normally accompany it as found in its native state. Thus, the peptides of this invention do not contain materials normally associated with their in situ environment, e.g., MHC I molecules on antigen presenting cells. Even where a protein has been isolated to a homogenous or dominant band, there are trace contaminants in the range of 5-10% of native protein which co-purify with the desired protein. Isolated peptides of this invention do not contain such endogenous co-purified protein.

The term "residue" refers to an amino acid or amino acid mimetic incorporated in an oligopeptide by an amide bond or amide bond mimetic.

## DESCRIPTION OF THE PREFERRED EMBODIMENTS

### I. HLA-A2.1 Motif

The present invention relates to the determination of allele-specific peptide motifs for human Class I MHC (sometimes referred to as HLA) allele subtypes, in

particular, peptide motifs recognized by HLA-A2.1 alleles. These motifs are then used to define T cell epitopes from any desired antigen, particularly those associated with human viral diseases, cancers or autoimmune diseases, for which the amino acid sequence of the potential antigen or autoantigen targets is known.

Epitopes on a number of potential target proteins can be identified in this manner. Examples of suitable antigens include prostate specific antigen (PSA), hepatitis B core and surface antigens (HBVc, HBVs) hepatitis C antigens, Epstein-Barr virus antigens, melanoma antigens (e.g., MAGE-1), human immunodeficiency virus (HIV) antigens, human papilloma virus (HPV) antigens, Lassa virus, mycobacterium tuberculosis (MT), p53, CEA, trypanosome surface antigen (TSA) and Her2/neu.

Peptides comprising the epitopes from these antigens are synthesized and then tested for their ability to bind to the appropriate MHC molecules in assays using, for example, purified class I molecules and radioiodinated peptides and/or cells expressing empty class I molecules by, for instance, immunofluorescent staining and flow microfluorometry, peptide-dependent class I assembly assays, and inhibition of CTL recognition by peptide competition. Those peptides that bind to the class I molecule are further evaluated for their ability to serve as targets for CTLs derived from infected or immunized individuals, as well as for their capacity to induce primary in vitro or in vivo CTL responses that can give rise to CTL populations capable of reacting with virally infected target cells or tumor cells as potential therapeutic agents.

The MHC class I antigens are encoded by the HLA-A, B, and C loci. HLA-A and B antigens are expressed at the cell surface at approximately equal densities, whereas the expression of HLA-C is significantly lower (perhaps as much as 10-fold lower). Each of these loci have a number of alleles. The peptide binding motifs of the invention are relatively specific for each allelic subtype.

For peptide-based vaccines, the peptides of the present invention preferably comprise a motif recognized by an MHC I molecule having a wide distribution in the human population. Since the MHC alleles occur at different frequencies within different ethnic groups and races, the choice of target MHC allele may depend upon the target population. Table 1 shows the frequency of various alleles at the HLA-A locus products among different races. For instance, the majority of the Caucasoid population can be covered by peptides which bind to four HLA-A allele subtypes, specifically HLA-



A2.1, A1, A3.2, and A24.1. Similarly, the majority of the Asian population is encompassed with the addition of peptides binding to a fifth allele HLA-A11.2.

TABLE 1

A Allele/Subtype	N(69)*	A(54)	C(502)
A1	10.1(7)	1.8(1)	27.4(138)
A2.1	11.5(8)	37.0(20)	39.8(199)
A2.2	10.1(7)	0	3.3(17)
A2.3	1.4(1)	5.5(3)	0.8(4)
A2.4	-	-	-
A2.5	-	-	-
A3.1	1.4(1)	0	0.2(0)
A3.2	5.7(4)	5.5(3)	21.5(108)
A11.1	0	5.5(3)	0
A11.2	5.7(4)	31.4(17)	8.7(44)
A11.3	0	3.7(2)	0
A23	4.3(3)	-	3.9(20)
A24	2.9(2)	27.7(15)	15.3(77)
A24.2	-	-	-
A24.3	-	-	-
A25	1.4(1)	-	6.9(35)
A26.1	4.3(3)	9.2(5)	5.9(30)
A26.2	7.2(5)	-	1.0(5)
A26V	-	3.7(2)	-
A28.1	10.1(7)	-	1.6(8)
A28.2	1.4(1)	-	7.5(38)
A29.1	1.4(1)	-	1.4(7)
A29.2	10.1(7)	1.8(1)	5.3(27)
A30.1	8.6(6)	-	4.9(25)
A30.2	1.4(1)	-	0.2(1)
A30.3	7.2(5)	-	3.9(20)
A31	4.3(3)	7.4(4)	6.9(35)
A32	2.8(2)	-	7.1(36)
Aw33.1	8.6(6)	-	2.5(13)
Aw33.2	2.8(2)	16.6(9)	1.2(6)
Aw34.1	1.4(1)	-	-
Aw34.2	14.5(10)	-	0.8(4)
Aw36	5.9(4)	-	-

Table compiled from B. DuPont, Immunobiology of HLA, Vol. I, Histocompatibility Testing 1987, Springer-Verlag, New York 1989.

5

\* N - negroid; A = Asian; C = Caucasoid. Numbers in parenthesis represent the number of individuals included in the analysis.

The nomenclature used to describe peptide compounds follows the

10

conventional practice wherein the amino group is presented to the left (the N-terminus) and the carboxyl group to the right (the C-terminus) of each amino acid residue. In the formulae representing selected specific embodiments of the present invention, the amino- and carboxyl-terminal groups, although not specifically shown, are in the form they would assume at physiologic pH values, unless otherwise specified. In the amino acid

structure formulae, each residue is generally represented by standard three letter or single letter designations. The L-form of an amino acid residue is represented by a capital single letter or a capital first letter of a three-letter symbol, and the D-form for those amino acids having D-forms is represented by a lower case single letter or a lower case three letter symbol. Glycine has no asymmetric carbon atom and is simply referred to as "Gly" or G.

The procedures used to identify peptides of the present invention generally follow the methods disclosed in Falk et al., Nature 351:290 (1991), which is incorporated herein by reference. Briefly, the methods involve large-scale isolation of MHC class I molecules, typically by immunoprecipitation or affinity chromatography, from the appropriate cell or cell line. Examples of other methods for isolation of the desired MHC molecule equally well known to the artisan include ion exchange chromatography, lectin chromatography, size exclusion, high performance ligand chromatography, and a combination of all of the above techniques.

In the typical case, immunoprecipitation is used to isolate the desired allele. A number of protocols can be used, depending upon the specificity of the antibodies used. For example, allele-specific mAb reagents can be used for the affinity purification of the HLA-A, HLA-B1, and HLA-C molecules. Several mAb reagents for the isolation of HLA-A molecules are available. The monoclonal BB7.2 is suitable for isolating HLA-A2 molecules. Affinity columns prepared with these mAbs using standard techniques are successfully used to purify the respective HLA-A allele products.

In addition to allele-specific mAbs, broadly reactive anti-HLA-A, B, C mAbs, such as W6/32 and B9.12.1, and one anti-HLA-B, C mAb, B1.23.2, could be used in alternative affinity purification protocols as described in previous applications.

The peptides bound to the peptide binding groove of the isolated MHC molecules are eluted typically using acid treatment. Peptides can also be dissociated from class I molecules by a variety of standard denaturing means, such as heat, pH, detergents, salts, chaotropic agents, or a combination thereof.

Peptide fractions are further separated from the MHC molecules by reversed-phase high performance liquid chromatography (HPLC) and sequenced.

Peptides can be separated by a variety of other standard means well known to the artisan, including filtration, ultrafiltration, electrophoresis, size chromatography, precipitation with specific antibodies, ion exchange chromatography, isoelectrofocusing, and the like.

Sequencing of the isolated peptides can be performed according to standard techniques such as Edman degradation (Hunkapiller, M.W., et al., Methods Enzymol. 91, 399 [1983]). Other methods suitable for sequencing include mass spectrometry sequencing of individual peptides as previously described (Hunt, et al., Science 225:1261 (1992), which is incorporated herein by reference). Amino acid sequencing of bulk heterogeneous peptides (e.g., pooled HPLC fractions) from different class I molecules typically reveals a characteristic sequence motif for each class I allele.

Definition of motifs specific for different class I alleles allows the identification of potential peptide epitopes from an antigenic protein whose amino acid sequence is known. Typically, identification of potential peptide epitopes is initially carried out using a computer to scan the amino acid sequence of a desired antigen for the presence of motifs. The epitopic sequences are then synthesized. The capacity to bind MHC Class molecules is measured in a variety of different ways. One means is a Class I molecule binding assay as described in the related applications, noted above. Other alternatives described in the literature include inhibition of antigen presentation (Sette, et al., J. Immunol. 141:3893 (1991), in vitro assembly assays (Townsend, et al., Cell 62:285 (1990), and FACS based assays using mutated cells, such as RMA.S (Melief, et al., Eur. J. Immunol. 21:2963 (1991)).

Next, peptides that test positive in the MHC class I binding assay are assayed for the ability of the peptides to induce specific CTL responses in vitro. For instance, antigen-presenting cells that have been incubated with a peptide can be assayed for the ability to induce CTL responses in responder cell populations. Antigen-presenting cells can be normal cells such as peripheral blood mononuclear cells or dendritic cells (Inaba, et al., J. Exp. Med. 166:182 (1987); Boog, Eur. J. Immunol. 18:219 [1988]).

Alternatively, mutant mammalian cell lines that are deficient in their ability to load class I molecules with internally processed peptides, such as the mouse cell lines RMA-S (Kärre, et al., Nature, 319:675 (1986); Ljunggren, et al., Eur. J. Immunol. 21:2963-2970 (1991)), and the human somatic T cell hybrid, T-2 (Cerundolo, et al., Nature 345:449-452 (1990)) and which have been transfected with the appropriate human class I genes are conveniently used, when peptide is added to them, to test for the capacity of the peptide to induce in vitro primary CTL responses. Other eukaryotic cell lines which could be used include various insect cell lines such as mosquito larvae (ATCC cell lines CCL 125, 126, 1660, 1591, 6585, 6586), silkworm (ATTC CRL 8851), armyworm

(ATCC CRL 1711), moth (ATCC CCL 80) and *Drosophila* cell lines such as a Schneider cell line (see Schneider *J. Embryol. Exp. Morphol.* 27:353-365 [1927]).

Peripheral blood lymphocytes are conveniently isolated following simple venipuncture or leukapheresis of normal donors or patients and used as the responder cell sources of CTL precursors. In one embodiment, the appropriate antigen-presenting cells are incubated with 10-100  $\mu$ M of peptide in serum-free media for 4 hours under appropriate culture conditions. The peptide-loaded antigen-presenting cells are then incubated with the responder cell populations *in vitro* for 7 to 10 days under optimized culture conditions. Positive CTL activation can be determined by assaying the cultures for the presence of CTLs that kill radiolabeled target cells, both specific peptide-pulsed targets as well as target cells expressing the endogenously processed form of the relevant virus or tumor antigen from which the peptide sequence was derived.

Specificity and MHC restriction of the CTL is determined by testing against different peptide target cells expressing appropriate or inappropriate human MHC class I. The peptides that test positive in the MHC binding assays and give rise to specific CTL responses are referred to herein as immunogenic peptides.

The immunogenic peptides can be prepared synthetically, or by recombinant DNA technology or from natural sources such as whole viruses or tumors. Although the peptide will preferably be substantially free of other naturally occurring host cell proteins and fragments thereof, in some embodiments the peptides can be synthetically conjugated to native fragments or particles.

The polypeptides or peptides can be a variety of lengths, either in their neutral (uncharged) forms or in forms which are salts, and either free of modifications such as glycosylation, side chain oxidation, or phosphorylation or containing these modifications, subject to the condition that the modification not destroy the biological activity of the polypeptides as herein described.

Desirably, the peptide will be as small as possible while still maintaining substantially all of the biological activity of the large peptide. When possible, it may be desirable to optimize peptides of the invention to a length of 9 or 10 amino acid residues, commensurate in size with endogenously processed viral peptides or tumor cell peptides that are bound to MHC class I molecules on the cell surface.

Peptides having the desired activity may be modified as necessary to provide certain desired attributes, e.g., improved pharmacological characteristics, while

increasing or at least retaining substantially all of the biological activity of the unmodified peptide to bind the desired MHC molecule and activate the appropriate T cell. For instance, the peptides may be subject to various changes, such as substitutions, either conservative or non-conservative, where such changes might provide for certain advantages in their use, such as improved MHC binding. By conservative substitutions is meant replacing an amino acid residue with another which is biologically and/or chemically similar, e.g., one hydrophobic residue for another, or one polar residue for another. The substitutions include combinations such as Gly, Ala; Val, Ile, Leu, Met; Asp, Glu; Asn, Gln; Ser, Thr; Lys, Arg; and Phe, Tyr. The effect of single amino acid substitutions may also be probed using D-amino acids. Such modifications may be made using well known peptide synthesis procedures, as described in e.g., Merrifield, Science 232:341-347 (1986), Barany and Merrifield, The Peptides, Gross and Meienhofer, eds. (N.Y., Academic Press), pp. 1-284 (1979); and Stewart and Young, Solid Phase Peptide Synthesis, (Rockford, Ill., Pierce), 2d Ed. (1984), incorporated by reference herein.

The peptides can also be modified by extending or decreasing the compound's amino acid sequence, e.g., by the addition or deletion of amino acids. The peptides or analogs of the invention can also be modified by altering the order or composition of certain residues, it being readily appreciated that certain amino acid residues essential for biological activity, e.g., those at critical contact sites or conserved residues, may generally not be altered without an adverse effect on biological activity. The non-critical amino acids need not be limited to those naturally occurring in proteins, such as L- $\alpha$ -amino acids, or their D-isomers, but may include non-natural amino acids as well, such as  $\beta$ - $\gamma$ - $\delta$ -amino acids, as well as many derivatives of L- $\alpha$ -amino acids.

Typically, a series of peptides with single amino acid substitutions are employed to determine the effect of electrostatic charge, hydrophobicity, etc. on binding. For instance, a series of positively charged (e.g., Lys or Arg) or negatively charged (e.g., Glu) amino acid substitutions are made along the length of the peptide revealing different patterns of sensitivity towards various MHC molecules and T cell receptors. In addition, multiple substitutions using small, relatively neutral moieties such as Ala, Gly, Pro, or similar residues may be employed. The substitutions may be homo-oligomers or hetero-oligomers. The number and types of residues which are substituted or added depend on the spacing necessary between essential contact points and certain functional attributes which are sought (e.g., hydrophobicity versus hydrophilicity). Increased binding affinity

for an MHC molecule or T cell receptor may also be achieved by such substitutions, compared to the affinity of the parent peptide. In any event, such substitutions should employ amino acid residues or other molecular fragments chosen to avoid, for example, steric and charge interference which might disrupt binding.

- 5                   Amino acid substitutions are typically of single residues. Substitutions, deletions, insertions or any combination thereof may be combined to arrive at a final peptide. Substitutional variants are those in which at least one residue of a peptide has been removed and a different residue inserted in its place. Such substitutions generally are made in accordance with the following Table 2 when it is desired to finely modulate
- 10 the characteristics of the peptide.

TABLE 2

<u>Original Residue</u>	<u>Exemplary Substitution</u>
Ala	Ser
Arg	Lys, His
Asn	Gln
Asp	Glu
Cys	Ser
Gln	Asn
Glu	Asp
Gly	Pro
His	Lys; Arg
Ile	Leu; Val
Leu	Ile; Val
Lys	Arg; His
Met	Leu; Ile
Phe	Tyr; Trp
Ser	Thr
Thr	Ser
Trp	Tyr; Phe
Tyr	Trp; Phe
Val	Ile; Leu

Pro

Gly

Substantial changes in function (e.g., affinity for MHC molecules or T cell receptors) are made by selecting substitutions that are less conservative than those in

5 Table 2, i.e., selecting residues that differ more significantly in their effect on maintaining (a) the structure of the peptide backbone in the area of the substitution, for example as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site or (c) the bulk of the side chain. The substitutions which in general are expected to produce the greatest changes in peptide properties will be those in which (a)

10 hydrophilic residue, e.g. seryl, is substituted for (or by) a hydrophobic residue, e.g. leucyl, isoleucyl, phenylalanyl, valyl or alanyl; (b) a residue having an electropositive side chain, e.g., lysyl, arginyl, or histidyl, is substituted for (or by) an electronegative residue, e.g. glutamyl or aspartyl; or (c) a residue having a bulky side chain, e.g. phenylalanine, is substituted for (or by) one not having a side chain, e.g., glycine.

15 The peptides may also comprise isosteres of two or more residues in the immunogenic peptide. An isostere as defined here is a sequence of two or more residues that can be substituted for a second sequence because the steric conformation of the first sequence fits a binding site specific for the second sequence. The term specifically includes peptide backbone modifications well known to those skilled in the art. Such

20 modifications include modifications of the amide nitrogen, the  $\alpha$ -carbon, amide carbonyl, complete replacement of the amide bond, extensions, deletions or backbone crosslinks. See, generally, Spatola, Chemistry and Biochemistry of Amino Acids, peptides and Proteins, Vol. VII (Weinstein ed., 1983).

25 Modifications of peptides with various amino acid mimetics or unnatural amino acids are particularly useful in increasing the stability of the peptide in vivo. Stability can be assayed in a number of ways. For instance, peptidases and various biological media, such as human plasma and serum, have been used to test stability. See, e.g., Verhoef et al., Eur. J. Drug Metab. Pharmacokin. 11:291-302 (1986). Half life of the peptides of the present invention is conveniently determined using a 25% human

30 serum (v/v) assay. The protocol is generally as follows. Pooled human serum (Type AB, non-heat inactivated) is delipidated by centrifugation before use. The serum is then diluted to 25% with RPMI tissue culture media and used to test peptide stability. At



predetermined time intervals a small amount of reaction solution is removed and added to either 6% aqueous trichloroacetic acid or ethanol. The cloudy reaction sample is cooled (4°C) for 15 minutes and then spun to pellet the precipitated serum proteins. The presence of the peptides is then determined by reversed-phase HPLC using stability-specific chromatography conditions.

The peptides of the present invention or analogs thereof which have CTL stimulating activity may be modified to provide desired attributes other than improved serum half life. For instance, the ability of the peptides to induce CTL activity can be enhanced by linkage to a sequence which contains at least one epitope that is capable of inducing a T helper cell response. Particularly preferred immunogenic peptides/T helper conjugates are linked by a spacer molecule. The spacer is typically comprised of relatively small, neutral molecules, such as amino acids or amino acid mimetics, which are substantially uncharged under physiological conditions. The spacers are typically selected from, e.g., Ala, Gly, or other neutral spacers of nonpolar amino acids or neutral polar amino acids. It will be understood that the optionally present spacer need not be comprised of the same residues and thus may be a hetero- or homo-oligomer. When present, the spacer will usually be at least one or two residues, more usually three to six residues. Alternatively, the CTL peptide may be linked to the T helper peptide without a spacer.

The immunogenic peptide may be linked to the T helper peptide either directly or via a spacer either at the amino or carboxy terminus of the CTL peptide. The amino terminus of either the immunogenic peptide or the T helper peptide may be acylated. Exemplary T helper peptides include tetanus toxoid 830-843, influenza 307-319, malaria circumsporozoite 382-398 and 378-389.

In some embodiments it may be desirable to include in the pharmaceutical compositions of the invention at least one component which primes CTL. Lipids have been identified as agents capable of priming CTL in vivo against viral antigens. For example, palmitic acid residues can be attached to the alpha and epsilon amino groups of a Lys residue and then linked, e.g., via one or more linking residues such as Gly, Gly-Gly-, Ser, Ser-Ser, or the like, to an immunogenic peptide. The lipidated peptide can then be injected directly in a micellar form, incorporated into a liposome or emulsified in an adjuvant, e.g., incomplete Freund's adjuvant. In a preferred embodiment a particularly effective immunogen comprises palmitic acid attached to alpha and epsilon amino groups

of Lys, which is attached via linkage, e.g., Ser-Ser, to the amino terminus of the immunogenic peptide.

As another example of lipid priming of CTL responses, *E. coli* lipoproteins, such as tripalmitoyl-S-glycerylcysteinylserine (P<sub>3</sub>CSS) can be used to prime virus specific CTL when covalently attached to an appropriate peptide. See, Deres et al., Nature 342:561-564 (1989), incorporated herein by reference. Peptides of the invention can be coupled to P<sub>3</sub>CSS, for example, and the lipopeptide administered to an individual to specifically prime a CTL response to the target antigen. Further, as the induction of neutralizing antibodies can also be primed with P<sub>3</sub>CSS conjugated to a peptide which displays an appropriate epitope, the two compositions can be combined to more effectively elicit both humoral and cell-mediated responses to infection.

In addition, additional amino acids can be added to the termini of a peptide to provide for ease of linking peptides one to another, for coupling to a carrier support, or larger peptide, for modifying the physical or chemical properties of the peptide or oligopeptide, or the like. Amino acids such as tyrosine, cysteine, lysine, glutamic or aspartic acid, or the like, can be introduced at the C- or N-terminus of the peptide or oligopeptide. Modification at the C terminus in some cases may alter binding characteristics of the peptide. In addition, the peptide or oligopeptide sequences can differ from the natural sequence by being modified by terminal-NH<sub>2</sub> acylation, e.g., by alkanoyl (C<sub>1</sub>-C<sub>20</sub>) or thioglycolyl acetylation, terminal-carboxyl amidation, e.g., ammonia, methylamine, etc. In some instances these modifications may provide sites for linking to a support or other molecule.

The peptides of the invention can be prepared in a wide variety of ways. Because of their relatively short size, the peptides can be synthesized in solution or on a solid support in accordance with conventional techniques. Various automatic synthesizers are commercially available and can be used in accordance with known protocols. See, for example, Stewart and Young, Solid Phase Peptide Synthesis, 2d. ed., Pierce Chemical Co. (1984), *supra*.

Alternatively, recombinant DNA technology may be employed wherein a nucleotide sequence which encodes an immunogenic peptide of interest is inserted into an expression vector, transformed or transfected into an appropriate host cell and cultivated under conditions suitable for expression. These procedures are generally known in the art, as described generally in Sambrook et al., Molecular Cloning, A Laboratory Manual,

Cold Spring Harbor Press, Cold Spring Harbor, New York (1982), which is incorporated herein by reference. Thus, fusion proteins which comprise one or more peptide sequences of the invention can be used to present the appropriate T cell epitope.

As the coding sequence for peptides of the length contemplated herein can be synthesized by chemical techniques, for example, the phosphotriester method of Matteucci et al., J. Am. Chem. Soc. 103:3185 (1981), modification can be made simply by substituting the appropriate base(s) for those encoding the native peptide sequence. The coding sequence can then be provided with appropriate linkers and ligated into expression vectors commonly available in the art, and the vectors used to transform suitable hosts to produce the desired fusion protein. A number of such vectors and suitable host systems are now available. For expression of the fusion proteins, the coding sequence will be provided with operably linked start and stop codons, promoter and terminator regions and usually a replication system to provide an expression vector for expression in the desired cellular host. For example, promoter sequences compatible with bacterial hosts are provided in plasmids containing convenient restriction sites for insertion of the desired coding sequence. The resulting expression vectors are transformed into suitable bacterial hosts. Of course, yeast or mammalian cell hosts may also be used, employing suitable vectors and control sequences.

The peptides of the present invention and pharmaceutical and vaccine compositions thereof are useful for administration to mammals, particularly humans, to treat and/or prevent viral infection and cancer. Examples of diseases which can be treated using the immunogenic peptides of the invention include prostate cancer, hepatitis B, hepatitis C, AIDS, renal carcinoma, cervical carcinoma, lymphoma, CMV and condyloma acuminatum.

For pharmaceutical compositions, the immunogenic peptides of the invention are administered to an individual already suffering from cancer or infected with the virus of interest. Those in the incubation phase or the acute phase of infection can be treated with the immunogenic peptides separately or in conjunction with other treatments, as appropriate. In therapeutic applications, compositions are administered to a patient in an amount sufficient to elicit an effective CTL response to the virus or tumor antigen and to cure or at least partially arrest symptoms and/or complications. An amount adequate to accomplish this is defined as "therapeutically effective dose." Amounts effective for this use will depend on, e.g., the peptide composition, the manner of administration, the stage

and severity of the disease being treated, the weight and general state of health of the patient, and the judgment of the prescribing physician, but generally range for the initial immunization (that is for therapeutic or prophylactic administration) from about 1.0  $\mu\text{g}$  to about 5000  $\mu\text{g}$  of peptide for a 70 kg patient, followed by boosting dosages of from about 1.0  $\mu\text{g}$  to about 1000  $\mu\text{g}$  of peptide pursuant to a boosting regimen over weeks to months depending upon the patient's response and condition by measuring specific CTL activity in the patient's blood. It must be kept in mind that the peptides and compositions of the present invention may generally be employed in serious disease states, that is, life-threatening or potentially life threatening situations. In such cases, in view of the minimization of extraneous substances and the relative nontoxic nature of the peptides, it is possible and may be felt desirable by the treating physician to administer substantial excesses of these peptide compositions.

For therapeutic use, administration should begin at the first sign of viral infection or the detection or surgical removal of tumors or shortly after diagnosis in the case of acute infection. This is followed by boosting doses until at least symptoms are substantially abated and for a period thereafter. In chronic infection, loading doses followed by boosting doses may be required.

Treatment of an infected individual with the compositions of the invention may hasten resolution of the infection in acutely infected individuals. For those individuals susceptible (or predisposed) to developing chronic infection the compositions are particularly useful in methods for preventing the evolution from acute to chronic infection. Where the susceptible individuals are identified prior to or during infection, for instance, as described herein, the composition can be targeted to them, minimizing need for administration to a larger population.

The peptide compositions can also be used for the treatment of chronic infection and to stimulate the immune system to eliminate virus-infected cells in carriers. It is important to provide an amount of immuno-potentiating peptide in a formulation and mode of administration sufficient to effectively stimulate a cytotoxic T cell response. Thus, for treatment of chronic infection, a representative dose is in the range of about 1.0  $\mu\text{g}$  to about 5000  $\mu\text{g}$ , preferably about 5  $\mu\text{g}$  to 1000  $\mu\text{g}$  for a 70 kg patient per dose. Immunizing doses followed by boosting doses at established intervals, e.g., from one to four weeks, may be required, possibly for a prolonged period of time to effectively immunize an individual. In the case of chronic infection, administration should continue

until at least clinical symptoms or laboratory tests indicate that the viral infection has been eliminated or substantially abated and for a period thereafter.

The pharmaceutical compositions for therapeutic treatment are intended for parenteral, topical, oral or local administration. Preferably, the pharmaceutical compositions are administered parenterally, e.g., intravenously, subcutaneously, intradermally, or intramuscularly. Thus, the invention provides compositions for parenteral administration which comprise a solution of the immunogenic peptides dissolved or suspended in an acceptable carrier, preferably an aqueous carrier. A variety of aqueous carriers may be used, e.g., water, buffered water, 0.8% saline, 0.3% glycine, hyaluronic acid and the like. These compositions may be sterilized by conventional, well known sterilization techniques, or may be sterile filtered. The resulting aqueous solutions may be packaged for use as is, or lyophilized, the lyophilized preparation being combined with a sterile solution prior to administration. The compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions, such as pH adjusting and buffering agents, tonicity adjusting agents, wetting agents and the like, for example, sodium acetate, sodium lactate, sodium chloride, potassium chloride, calcium chloride, sorbitan monolaurate, triethanolamine oleate, etc.

The concentration of CTL stimulatory peptides of the invention in the pharmaceutical formulations can vary widely, i.e., from less than about 0.1%, usually at or at least about 2% to as much as 20% to 50% or more by weight, and will be selected primarily by fluid volumes, viscosities, etc., in accordance with the particular mode of administration selected.

The peptides of the invention may also be administered via liposomes, which serve to target the peptides to a particular tissue, such as lymphoid tissue, or targeted selectively to infected cells, as well as increase the half-life of the peptide composition. Liposomes include emulsions, foams, micelles, insoluble monolayers, liquid crystals, phospholipid dispersions, lamellar layers and the like. In these preparations the peptide to be delivered is incorporated as part of a liposome, alone or in conjunction with a molecule which binds to, e.g., a receptor prevalent among lymphoid cells, such as monoclonal antibodies which bind to the CD45 antigen, or with other therapeutic or immunogenic compositions. Thus, liposomes either filled or decorated with a desired peptide of the invention can be directed to the site of lymphoid cells, where

the liposomes then deliver the selected therapeutic/immunogenic peptide compositions. Liposomes for use in the invention are formed from standard vesicle-forming lipids, which generally include neutral and negatively charged phospholipids and a sterol, such as cholesterol. The selection of lipids is generally guided by consideration of, e.g., liposome size, acid lability and stability of the liposomes in the blood stream. A variety of methods are available for preparing liposomes, as described in, e.g., Szoka et al., Ann. Rev. Biophys. Bioeng. 9:467 (1980), U.S. Patent Nos. 4,235,871, 4,501,728, 4,837,028, and 5,019,369, incorporated herein by reference.

For targeting to the immune cells, a ligand to be incorporated into the liposome can include, e.g., antibodies or fragments thereof specific for cell surface determinants of the desired immune system cells. A liposome suspension containing a peptide may be administered intravenously, locally, topically, etc. in a dose which varies according to, inter alia, the manner of administration, the peptide being delivered, and the stage of the disease being treated.

For solid compositions, conventional nontoxic solid carriers may be used which include, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharin, talcum, cellulose, glucose, sucrose, magnesium carbonate, and the like. For oral administration, a pharmaceutically acceptable nontoxic composition is formed by incorporating any of the normally employed excipients, such as those carriers previously listed, and generally 10-95% of active ingredient, that is, one or more peptides of the invention, and more preferably at a concentration of 25%-75%.

For aerosol administration, the immunogenic peptides are preferably supplied in finely divided form along with a surfactant and propellant. Typical percentages of peptides are 0.01%-20% by weight, preferably 1%-10%. The surfactant must, of course, be nontoxic, and preferably soluble in the propellant. Representative of such agents are the esters or partial esters of fatty acids containing from 6 to 22 carbon atoms, such as caproic, octanoic, lauric, palmitic, stearic, linoleic, linolenic, olesteric and oleic acids with an aliphatic polyhydric alcohol or its cyclic anhydride. Mixed esters, such as mixed or natural glycerides may be employed. The surfactant may constitute 0.1%-20% by weight of the composition, preferably 0.25-5%. The balance of the composition is ordinarily propellant. A carrier can also be included, as desired, as with, e.g., lecithin for intranasal delivery.

In another aspect the present invention is directed to vaccines which contain as an active ingredient an immunogenically effective amount of an immunogenic peptide as described herein. The peptide(s) may be introduced into a host, including humans, linked to its own carrier or as a homopolymer or heteropolymer of active peptide units. Such a polymer has the advantage of increased immunological reaction and, where different peptides are used to make up the polymer, the additional ability to induce antibodies and/or CTLs that react with different antigenic determinants of the virus or tumor cells. Useful carriers are well known in the art, and include, e.g., thyroglobulin, albumins such as human serum albumin, tetanus toxoid, polyamino acids such as poly(lysine:glutamic acid), influenza, hepatitis B virus core protein, hepatitis B virus recombinant vaccine and the like. The vaccines can also contain a physiologically tolerable (acceptable) diluent such as water, phosphate buffered saline, or saline, and further typically include an adjuvant. Adjuvants such as incomplete Freund's adjuvant, aluminum phosphate, aluminum hydroxide, or alum are materials well known in the art. And, as mentioned above, CTL responses can be primed by conjugating peptides of the invention to lipids, such as P3CSS. Upon immunization with a peptide composition as described herein, via injection, aerosol, oral, transdermal or other route, the immune system of the host responds to the vaccine by producing large amounts of CTLs specific for the desired antigen, and the host becomes at least partially immune to later infection, or resistant to developing chronic infection.

Vaccine compositions containing the peptides of the invention are administered to a patient susceptible to or otherwise at risk of viral infection or cancer to elicit an immune response against the antigen and thus enhance the patient's own immune response capabilities. Such an amount is defined to be an "immunogenically effective dose." In this use, the precise amounts again depend on the patient's state of health and weight, the mode of administration, the nature of the formulation, etc., but generally range from about 1.0  $\mu\text{g}$  to about 5000  $\mu\text{g}$  per 70 kilogram patient, more commonly from about 10  $\mu\text{g}$  to about 500  $\mu\text{g}$  mg per 70 kg of body weight.

In some instances it may be desirable to combine the peptide vaccines of the invention with vaccines which induce neutralizing antibody responses to the virus of interest, particularly to viral envelope antigens.

For therapeutic or immunization purposes, nucleic acids encoding one or more of the peptides of the invention can also be administered to the patient. A number

of methods are conveniently used to deliver the nucleic acids to the patient. For instance, the nucleic acid can be delivered directly, as “naked DNA”. This approach is described, for instance, in Wolff *et. al.*, *Science* 247: 1465-1468 (1990) as well as U.S. Patent Nos. 5,580,859 and 5,589,466. The nucleic acids can also be administered using ballistic  
5 delivery as described, for instance, in U.S. Patent No. 5,204,253. Particles comprised solely of DNA can be administered. Alternatively, DNA can be adhered to particles, such as gold particles. The nucleic acids can also be delivered complexed to cationic compounds, such as cationic lipids. Lipid-mediated gene delivery methods are described, for instance, in WO 96/18372; WO 93/24640; Mannino and Gould-Fogerite (1988)  
10 *BioTechniques* 6(7): 682-691; Rose U.S. Pat No. 5,279,833; WO 91/06309; and Felgner *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84: 7413-7414. The peptides of the invention can also be expressed by attenuated viral hosts, such as vaccinia or fowlpox. This approach involves the use of vaccinia virus as a vector to express nucleotide sequences that encode the peptides of the invention. Upon introduction into an acutely or  
15 chronically infected host or into a noninfected host, the recombinant vaccinia virus expresses the immunogenic peptide, and thereby elicits a host CTL response. Vaccinia vectors and methods useful in immunization protocols are described in, e.g., U.S. Patent No. 4,722,848, incorporated herein by reference. Another vector is BCG (Bacille Calmette Guerin). BCG vectors are described in Stover *et al.* (*Nature* 351:456-460  
20 (1991)) which is incorporated herein by reference. A wide variety of other vectors useful for therapeutic administration or immunization of the peptides of the invention, e.g., Salmonella typhi vectors and the like, will be apparent to those skilled in the art from the description herein.

A preferred means of administering nucleic acids encoding the peptides of  
25 the invention uses minigene constructs encoding multiple epitopes of the invention. To create a DNA sequence encoding the selected CTL epitopes (minigene) for expression in human cells, the amino acid sequences of the epitopes are reverse translated. A human codon usage table is used to guide the codon choice for each amino acid. These epitope-encoding DNA sequences are directly adjoined, creating a continuous polypeptide  
30 sequence. To optimize expression and/or immunogenicity, additional elements can be incorporated into the minigene design. Examples of amino acid sequence that could be reverse translated and included in the minigene sequence include: helper T lymphocyte epitopes, a leader (signal) sequence, and an endoplasmic reticulum retention signal. In



addition, MHC presentation of CTL epitopes may be improved by including synthetic (e.g. poly-alanine) or naturally-occurring flanking sequences adjacent to the CTL epitopes.

The minigene sequence is converted to DNA by assembling  
5 oligonucleotides that encode the plus and minus strands of the minigene. Overlapping oligonucleotides (30-100 bases long) are synthesized, phosphorylated, purified and annealed under appropriate conditions using well known techniques. The ends of the oligonucleotides are joined using T4 DNA ligase. This synthetic minigene, encoding the CTL epitope polypeptide, can then be cloned into a desired expression vector.

10 Standard regulatory sequences well known to those of skill in the art are included in the vector to ensure expression in the target cells. Several vector elements are required: a promoter with a down-stream cloning site for minigene insertion; a polyadenylation signal for efficient transcription termination; an *E. coli* origin of replication; and an *E. coli* selectable marker (e.g. ampicillin or kanamycin resistance).

15 Numerous promoters can be used for this purpose, e.g., the human cytomegalovirus (hCMV) promoter. See, U.S. Patent Nos. 5,580,859 and 5,589,466 for other suitable promoter sequences.

Additional vector modifications may be desired to optimize minigene expression and immunogenicity. In some cases, introns are required for efficient gene  
20 expression, and one or more synthetic or naturally-occurring introns could be incorporated into the transcribed region of the minigene. The inclusion of mRNA stabilization sequences can also be considered for increasing minigene expression. It has recently been proposed that immunostimulatory sequences (ISSs or CpGs) play a role in the immunogenicity of DNA vaccines. These sequences could be included in the vector,  
25 outside the minigene coding sequence, if found to enhance immunogenicity.

In some embodiments, a bioisotonic expression vector, to allow production of the minigene-encoded epitopes and a second protein included to enhance or decrease immunogenicity can be used. Examples of proteins or polypeptides that could  
30 beneficially enhance the immune response if co-expressed include cytokines (e.g., IL2, IL12, GM-CSF), cytokine-inducing molecules (e.g. LeIF) or costimulatory molecules. Helper (HTL) epitopes could be joined to intracellular targeting signals and expressed separately from the CTL epitopes. This would allow direction of the HTL epitopes to a cell compartment different than the CTL epitopes. If required, this could facilitate more

efficient entry of HTL epitopes into the MHC class II pathway, thereby improving CTL induction. In contrast to CTL induction, specifically decreasing the immune response by co-expression of immunosuppressive molecules (e.g. TGF- $\beta$ ) may be beneficial in certain diseases.

5                   Once an expression vector is selected, the minigene is cloned into the polylinker region downstream of the promoter. This plasmid is transformed into an appropriate *E. coli* strain, and DNA is prepared using standard techniques. The orientation and DNA sequence of the minigene, as well as all other elements included in the vector, are confirmed using restriction mapping and DNA sequence analysis.

10       Bacterial cells harboring the correct plasmid can be stored as a master cell bank and a working cell bank.

                  Therapeutic quantities of plasmid DNA are produced by fermentation in *E. coli*, followed by purification. Aliquots from the working cell bank are used to inoculate fermentation medium (such as Terrific Broth), and grown to saturation in shaker flasks or

15       a bioreactor according to well known techniques. Plasmid DNA can be purified using standard bioseparation technologies such as solid phase anion-exchange resins supplied by Quiagen. If required, supercoiled DNA can be isolated from the open circular and linear forms using gel electrophoresis or other methods.

                  Purified plasmid DNA can be prepared for injection using a variety of

20       formulations. The simplest of these is reconstitution of lyophilized DNA in sterile phosphate-buffer saline (PBS). A variety of methods have been described, and new techniques may become available. As noted above, nucleic acids are conveniently formulated with cationic lipids. In addition, glycolipids, fusogenic liposomes, peptides and compounds referred to collectively as protective, interactive, non-condensing (PINC)

25       could also be complexed to purified plasmid DNA to influence variables such as stability, intramuscular dispersion, or trafficking to specific organs or cell types.

                  Target cell sensitization can be used as a functional assay for expression and MHC class I presentation of minigene-encoded CTL epitopes. The plasmid DNA is introduced into a mammalian cell line that is suitable as a target for standard CTL

30       chromium release assays. The transfection method used will be dependent on the final formulation. Electroporation can be used for "naked" DNA, whereas cationic lipids allow direct in vitro transfection. A plasmid expressing green fluorescent protein (GFP) can be co-transfected to allow enrichment of transfected cells using fluorescence activated cell

sorting (FACS). These cells are then chromium-51 labeled and used as target cells for epitope-specific CTL lines. Cytolysis, detected by <sup>51</sup>Cr release, indicates production of MHC presentation of minigene-encoded CTL epitopes.

*In vivo* immunogenicity is a second approach for functional testing of minigene DNA formulations. Transgenic mice expressing appropriate human MHC molecules are immunized with the DNA product. The dose and route of administration are formulation dependent (e.g. IM for DNA in PBS, IP for lipid-complexed DNA). Twenty-one days after immunization, splenocytes are harvested and restimulated for 1 week in the presence of peptides encoding each epitope being tested. These effector cells (CTLs) are assayed for cytolysis of peptide-loaded, chromium-51 labeled target cells using standard techniques. Lysis of target cells sensitized by MHC loading of peptides corresponding to minigene-encoded epitopes demonstrates DNA vaccine function for *in vivo* induction of CTLs.

Antigenic peptides may be used to elicit CTL ex vivo, as well. The resulting CTL, can be used to treat chronic infections (viral or bacterial) or tumors in patients that do not respond to other conventional forms of therapy, or will not respond to a peptide vaccine approach of therapy. Ex vivo CTL responses to a particular pathogen (infectious agent or tumor antigen) are induced by incubating in tissue culture the patient's CTL precursor cells (CTLp) together with a source of antigen-presenting cells (APC) and the appropriate immunogenic peptide. After an appropriate incubation time (typically 1-4 weeks), in which the CTLp are activated and mature and expand into effector CTL, the cells are infused back into the patient, where they will destroy their specific target cell (an infected cell or a tumor cell).

The peptides may also find use as diagnostic reagents. For example, a peptide of the invention may be used to determine the susceptibility of a particular individual to a treatment regimen which employs the peptide or related peptides, and thus may be helpful in modifying an existing treatment protocol or in determining a prognosis for an affected individual. In addition, the peptides may also be used to predict which individuals will be at substantial risk for developing chronic infection.

The following example is offered by way of illustration, not by way of limitation.

## EXAMPLE 1

Class I antigen isolation was carried out as described in the related applications, noted above. Naturally processed peptides were then isolated and sequenced as described there. An allele-specific motif and algorithms were determined  
5 and quantitative binding assays were carried out.

Using the motifs identified above for the HLA-A2.1 allele amino acid sequences from a number of antigens were analyzed for the presence of these motifs. Table 3 provides the results of these searches. The letter "J" represents norleucine.

The above examples are provided to illustrate the invention but not to limit  
10 its scope. Other variants of the invention will be readily apparent to one of ordinary skill in the art and are encompassed by the appended claims. All publications, patents, and patent applications cited herein are hereby incorporated by reference.

Table 3

<u>Peptide</u>	<u>AA</u>	<u>Sequence</u>	<u>Source</u>	<u>A*0201</u>	<u>SEQ ID NO:</u>
17.0317	9	LQIGNIISI	Flu.24	0.0130	1
38.0103	9	NLSLSCHAA	CEA.432	0.0110	2
1233.11	9	YLSGANLNV	CEA.605V9	0.0690	3
1295.03	9	SMPPPGTRV	p53.149M2	0.0290	4
1295.04	9	SLPPPGTRV	p53.149L2	0.0410	5
1317.24	9	KTCPVQLWV	p53.139	0.0069	6
1323.02	9	KLLPENNVV	p53.24V9	0.0130	7
1323.04	9	ALNKMFBQV	p53.129B7V9	0.0260	8
1323.06	9	KLBPVQLWV	p53.139L2B3	0.1100	9
1323.08	9	BLTIHYNVYV	p53.229B1L2V9	0.0430	10
1323.18	10	LLPPQHLIRV	p53.188L2	0.0061	11
1323.29	11	YMCNSSCMGGM	p53.236	0.0075	12
1323.31	11	YLCNSSCMGGV	p53.236L2V11	0.2300	13
1323.34	11	KLYQGSYGFRV	p53.101L2V11	0.0620	14
1324.07	9	CQLAKTCPV	p53.135	0.0240	15
1325.01	9	RLPEAAPPV	p53.65L2	0.0640	16
1325.02	9	GLAPPQHLV	p53.187V9	0.0130	17
1325.04	9	KMAELVHFL	MAGE3.112M2	0.2100	18

<u>Peptide</u>	<u>AA</u>	<u>Sequence</u>	<u>Source</u>	<u>A*0201</u>	<u>SEQ ID NO:</u>
1325.05	9	KLAELVHFL	MAGE3.112L2	0.2500	19
1326.01	9	CLLAKTCPV	p53.135L2	0.0400	20
1326.02	9	KLSQHMTEV	p53.164L2	0.0410	21
1326.04	9	ELAPVVAPV	p53.68L2V9	0.0860	22
1326.06	10	QLAKTCPVQV	p53.136	0.0320	23
1326.08	9	HLTEVVRRV	p53.168L2	0.0180	24
1329.01	11	KTYQGSYGFR		0.0028	25
1329.03	10	VVVPYEPPEV	p53.216	0.0081	26
1329.14	9	BQLAKTBPV	p53.135B1B7	0.0490	27
1329.15	9	BLLAKTBPV	p53.135B1L2B7	0.1100	28
1330.01	9	QIIIGYVIGT	CEA.78	0.0160	29
1330.02	9	QLIGYVIGV	CEA.78L2V9	0.5300	30
1330.05	9	YVCGIQNSV	CEA.569	0.0510	31
1330.06	9	YLCGIQNSV	CEA.569L2	0.1000	32
1330.07	9	ATVGIMIGV	CEA.687	0.1400	33
1330.08	9	ALVGIMIGV	CEA.687L2	0.5000	34
1330.09	10	VLYGPDDPTI	CEA.411	0.0170	35
1330.10	10	VLYGPDDPTV	CEA.411V10	0.0310	36
1331.02	9	DLMLSPDDV	p53.42V9		37
1331.03	9	ALMLSPDDI	p53.42A1		38
1331.04	9	ALMLSPDDV	p53.42A1V9		39
1331.05	9	DLMLSPADI	p53.42A7		40
1331.06	9	DLMLSPADV	p53.42A7V9		41
1331.07	9	DLMLSPDAI	p53.42A8		42
1331.08	9	DLMLSPDAV	p53.42A8V9		43
38.0007	9	AILTFGSFV	KSHV.89	0.0850	44
38.0009	9	HLRDFALAV	KSHV.106	0.0183	45
38.0015	9	ALLGSIALL	KSHV.155	0.0470	46
38.0018	9	ALLATILAA	KSHV.161	0.0490	47
38.0019	9	LLATILAAV	KSHV.162	0.1600	48
38.0022	9	RLFADELAA	KSHV.14	0.0150	49

<u>Peptide</u>	<u>AA</u>	<u>Sequence</u>	<u>Source</u>	<u>A*0201</u>	<u>SEQ ID NO:</u>
38.0024	9	YLSKCTLAV	KSHV.65	0.2000	50
38.0026	9	LVYHIYSKI	KSHV.153	0.0457	51
38.0029	9	SMYLCILSA	KSHV.208	0.0250	52
38.0030	9	YLCILSALV	KSHV.210	0.3500	53
38.0033	9	VMFSYLQSL	KSHV.268	0.5000	54
38.0035	9	RLHVEAYSA	KSHV.285	0.0270	55
38.0039	9	GLQTLGAFV	KSHV.98	0.0110	56
38.0040	9	FVEEQMTWA	KSHV.105	0.0380	57
38.0041	9	QMTWAQTVV	KSHV.109	0.0110	58
38.0042	9	IILDTAIFV	KSHV.130	0.6800	59
38.0043	9	AIFVCNAFV	KSHV.135	0.0910	60
38.0046	9	AMGNRLVEA	KSHV.172	0.0200	61
38.0047	9	RLVEACNLL	KSHV.176	0.0180	62
38.0059	9	TLSIVTFSL	KSHV.198	0.2200	63
38.0063	9	KLSVLLLEV	KSHV.292	0.1400	64
38.0064	9	LLLEVNRSV	KSHV.296	0.0270	65
38.0068	9	FVSSPTLPV	KSHV.78	0.0350	66
38.0070	9	AMLVLLAEI	KSHV.281	0.0820	67
38.0075	9	QMARLAWEA	KSHV.1116	0.0990	68
38.0131	10	VLAIEGIFMA	KSHV.10	0.0730	69
38.0132	10	YLYHPLLSP	KSHV.27	0.1400	70
38.0134	10	SLFEAMLANV	KSHV.49	0.9500	71
38.0135	10	STTGINQLGL	KSHV.62	0.0710	72
38.0137	10	LAILTFGSFV	KSHV.88	0.0160	73
38.0139	10	ALLGSIALLA	KSHV.155	0.0360	74
38.0141	10	ALLATILAAV	KSHV.161	0.1100	75
38.0142	10	LLATILAAVA	KSHV.162	0.0110	76
38.0143	10	RLFADELAAL	KSHV.14	0.1800	77
38.0148	10	YLSKCTLAVL	KSHV.65	0.0300	78
38.0150	10	LLVYHIYSKI	KSHV.152	0.0130	79
38.0151	10	SMYLCILSAL	KSHV.208	0.0360	80

<u>Peptide</u>	<u>AA</u>	<u>Sequence</u>	<u>Source</u>	<u>A*0201</u>	<u>SEQ ID NO:</u>
38.0153	10	HLHRQMLSFV	KSHV.68	0.0160	81
38.0163	10	LLCGKTGAFL	KSHV.167	0.0100	82
38.0164	10	ETLSIVTFSL	KSHV.197	0.0180	83
39.0063	9	VMCTYSPPL	mp53.119	1.4000	84
39.0065	9	KLFCQLAKT	mp53.129	0.0160	85
39.0067	9	ATPPAGSRV	mp53.146	0.0130	86
39.0133	10	FLQSGTAKSV	mp53.110	0.0180	87
39.0169	10	CMDRGLTVFV	KSHV.311	0.0120	88
39.0170	10	VLLNWWRWRL	KSHV.327	0.1500	89
40.0070	9	GVFTGLTHI	HCV.1565	0.0110	90
40.0072	9	QMWKCLIRL	HCV.1611	0.0620	91
40.0074	9	IMTCMSADL	HCV.1650	0.0121	92
40.0076	9	ALAAAYCLST	HCV.1674	0.2500	93
40.0080	9	VLSGKPAII	HCV.1692	0.0150	94
40.0082	9	FISGIQYLA	HCV.1773	0.1000	95
40.0134	10	YIMTCMSADL	HCV.1649	0.0300	96
40.0137	10	AIASLMAFTA	HCV.1791	0.0580	97
40.0138	10	GLAGAAIGSV	HCV.1838	0.0320	98
41.0058	8	MIGVLVGV	CEA.692	0.0120	99
41.0061	9	VLPLAYISL	TRP1	0.0110	100
41.0062	9	SLGCIFFPL	TRP1	0.9700	101
41.0063	9	PLAYISLFL	TRP1	0.0220	102
41.0065	9	LMLFYQVWA	TRP1	0.0270	103
41.0071	9	NISIYNYFV	TRP1	0.2300	104
41.0072	9	NISVYNYFV	TRP1	0.0600	105
41.0075	9	FVWTHYYSV	TRP1	1.5000	106
41.0077	9	FLTWHRYHL	TRP1	0.5500	107
41.0078	9	LTWHRYHLL	TRP1	0.1600	108
41.0082	9	MLQEPSFSL	TRP1	0.6900	109
41.0083	9	SLPYWNFAT	TRP1	0.0110	110
41.0088	9	RLPEPQDVA	TRP1	0.0180	111

<u>Peptide</u>	<u>AA</u>	<u>Sequence</u>	<u>Source</u>	<u>A*0201</u>	<u>SEQ ID NO:</u>
41.0090	9	VTQCLEVRV	TRP1	0.0160	112
41.0096	9	LLHTFTDAV	TRP1	0.2700	113
41.0100	9	NMVPFWPPV	TRP1	0.6200	114
41.0104	9	AVVGALLLV	TRP1	0.0210	115
41.0105	9	AVVAALLLV	TRP1	0.0390	116
41.0108	9	LLVAAIFGV	TRP1	1.9000	117
41.0112	9	SMDEANQPL	TRP1	0.0770	118
41.0114	9	VLPLAYISV	TRP1	0.1100	119
41.0115	9	SLGCIFFPV	TRP1	3.2000	120
41.0116	9	PLAYISLFV	TRP1	0.0310	121
41.0117	9	LLLFQQARV	TRP1	0.1100	122
41.0118	9	LMLFYQVWV	TRP1	2.4000	123
41.0119	9	LLPSSGPGV	TRP1	0.3700	124
41.0121	9	NLSIYNYFV	TRP1	0.9700	125
41.0122	9	NLSVYNYFV	TRP1	0.8700	126
41.0123	9	FLWTHYYSV	TRP1	5.6000	127
41.0124	9	SLKKTFLGV	TRP1	0.0224	128
41.0125	9	FLTWHRYHV	TRP1	0.3800	129
41.0129	9	MLQEFSFSV	TRP1	1.6000	130
41.0130	9	SLPYWNFAV	TRP1	0.5700	131
41.0131	9	ALGKNVCDV	TRP1	0.0160	132
41.0132	9	SLISPNSV	TRP1	0.1300	133
41.0133	9	SLFSQWRVV	TRP1	0.0740	134
41.0134	9	TLGTLCNSV	TRP1	0.0330	135
41.0136	9	RLPEPQDVV	TRP1	0.1000	136
41.0137	9	VLQCLEVRV	TRP1	0.0360	137
41.0138	9	SLNSFRNTV	TRP1	0.0140	138
41.0139	9	SLDSFRNTV	TRP1	0.0440	139
41.0141	9	FLNGTGGQV	TRP1	0.0220	140
41.0142	9	VLLHTFTDV	TRP1	0.0180	141
41.0145	9	ALVGALLLV	TRP1	0.2600	142



<u>Peptide</u>	<u>AA</u>	<u>Sequence</u>	<u>Source</u>	<u>A*0201</u>	<u>SEQ ID NO:</u>
41.0146	9	ALVAALLLV	TRP1	0.5800	143
41.0147	9	LLVALIFGV	TRP1	1.0000	144
41.0148	9	YLIRARRSV	TRP1	0.0170	145
41.0149	9	SMDEANQPV	TRP1	0.1600	146
41.0151	10	SLGCIFFPLL	TRP1	0.1800	147
41.0157	10	GMCCPDLSPV	TRP1	0.0950	148
41.0160	10	AACNQKILTV	TRP1	0.0120	149
41.0162	10	FLTWHRYHLL	TRP1	0.0830	150
41.0166	10	SLHNLAHLFL	TRP1	0.3900	151
41.0174	10	LLLVAAIFGV	TRP1	0.3000	152
41.0177	10	LLVAAIFGVA	TRP1	0.0820	153
41.0178	10	ALIFGTASYL	TRP1	0.0230	154
41.0180	10	SMDEANQPLL	TRP1	0.0250	155
41.0181	10	LLTDQYQCYA	TRP1	0.0320	156
41.0183	10	SLGCIFFPLV	TRP1	0.3200	157
41.0186	10	FLMLFYQVWV	TRP1	0.8100	158
41.0189	10	ALCDQRV LIV	TRP1	0.0530	159
41.0190	10	ALCNQKILTV	TRP1	0.0770	160
41.0191	10	FLTWHRYHLV	TRP1	0.0510	161
41.0197	10	SLHNLAHLFV	TRP1	0.5000	162
41.0198	10	NLAHLFLNGV	TRP1	0.4100	163
41.0199	10	NMVPFWPPVV	TRP1	0.2800	164
41.0201	10	ILVVAALLLV	TRP1	0.0190	165
41.0203	10	LLVALIFGTV	TRP1	0.1200	166
41.0205	10	ALIFGTASYV	TRP1	0.0900	167
41.0206	10	SMDEANQPLV	TRP1	0.0350	168
41.0207	10	LLTDQYQCYV	TRP1	0.2100	169
41.0212	11	LLIQNIIQN DT	CEA.107	0.0140	170
41.0214	11	IIQN DTGFYTL	CEA.112	0.0130	171
41.0221	11	TLFNVTRNDTA	CEA.201	0.0110	172
41.0235	11	LTLLSVTRNDV	CEA.378	0.0150	173

<u>Peptide</u>	<u>AA</u>	<u>Sequence</u>	<u>Source</u>	<u>A*0201</u>	<u>SEQ ID NO:</u>
41.0243	11	GLYTCQANNSA	CEA.473	0.0290	174
41.0268	11	ATVGIMIGVLV	CEA.687	0.0160	175
44.0075	11	GLVPPQHLIRV	mp53.184.V3	0.0370	176
44.0087	11	GLAPPVHLIRV	mp53.184.V6	0.0330	177
44.0092	11	GLAPPEHLIRV	mp53.184.E6	0.1600	178
1227.10	9	ILIGVLVGV	CEA.691.L2	0.2300	179
1234.26	10	YLIMVKCWMV	Her2/neu.952.L2	0.3800	180
V10					
1295.06	9	LLGRDSFEV	mp53.261	0.2000	181
1319.01	9	FMYSDFHFI	Flu.RRP2.446	0.4400	182
1319.06	9	NMLSTVLGV	Flu.RRP2.446	0.1700	183
1319.14	9	SLENFRAYV	Flu.RRP2.446	0.0430	184
1325.06		KMAELVHFV	Mage3.112	0.1900	185
1325.07		KLAELVHFV	Mage3.112	0.3500	186
1334.01		VLIQRNPQV	Her2/neu.153.V9	0.0910	187
1334.02		VLLGVVFGV	Her2/neu.665.L2	2.1000	188
V9					
1334.03		SLISAVVGV	Her2/neu.653.L2	0.7000	189
V9					
1334.04		YMIMVKBWMI	Her2/neu.952.B7	0.2700	190
1334.05		YLIMVKBWMV	Her2/neu.952.L2	0.6900	191
B7V10					
1334.06		KLWEELSVV	Mage3.220.L2V	0.4500	192
9					
1334.08		AMBRWGLLV	Her2/neu.5.M2B	0.1400	193
3V9					
1345.01	9	IIGVLVGV	CEA.691.J2	0.0570	194
1345.02	9	ATVGIIIGV	CEA.687.J6	0.1595	195
1345.03	9	SJPPPGTRV	p53.149.J2	0.0545	196
1345.04	10	LVFGIELJEV	MAGE3.160.J8	0.7650	197
918.12	8	ILGFVFTL	Flu.M1.59	0.7900	198

<u>Peptide</u>	<u>AA</u>	<u>Sequence</u>	<u>Source</u>	A*0201	<u>SEQ ID NO:</u>
1095.22	9	KIFGSLAFL	Her2/neu		199
1090.01	10	YLQLVFGIEV	MAGE2		200
1126.01	9	MMNDQLMFL	PSM		201
1126.02	10	ALVLAGGFFL	PSM		202
1126.03	9	WLCAGALVL	PSM		203
1126.05	9	MVFELANSI	PSM		204
1126.06	10	RMMNDQLMFL	PSM		205
1126.09	9	LVLAGGFFL	PSM		206
1126.10	9	VLAGGFFLL	PSM		207
1126.12	9	LLHETDSAV	PSM		208
1126.14	9	LMYSLVHNL	PSM		209
1126.16	10	QLMFLERAFL	PSM		210
1126.17	9	LMFLERAFL	PSM		211
1126.20	10	KLGSNDFEV	PSM		212
1129.01	10	LLQERGVAYI	PSM		213
1129.04	10	GMPEGDLVYV	PSM		214
1129.05	10	FLDELKAENI	PSM		215
1129.08	9	ALFDIESKV	PSM		216
1129.10	10	GLPSIPVHPI	PSM		217

## II. Non-HLA-A2 Motifs

The present invention also relates to the determination of allele-specific peptide motifs for human Class I MHC (sometimes referred to as HLA) allele subtypes.

- 5 These motifs are then used to define T cell epitopes from any desired antigen, particularly those associated with human viral diseases, cancers or autoimmune diseases, for which the amino acid sequence of the potential antigen or autoantigen targets is known.

- Epitopes on a number of potential target proteins can be identified in this manner. Examples of suitable antigens include prostate specific antigen (PSA), hepatitis
- 10 B core and surface antigens (HBVc, HBVs) hepatitis C antigens, Epstein-Barr virus antigens, melanoma antigens (e.g., MAGE-1), human immunodeficiency virus (HIV) antigens and human papilloma virus (HPV) antigens, Lassa virus, mycobacterium tuberculosis (MT), p53, CEA, and Her2/neu.

Peptides comprising the epitopes from these antigens are synthesized and then tested for their ability to bind to the appropriate MHC molecules in assays using, for example, purified class I molecules and radioiodinated peptides and/or cells expressing empty class I molecules by, for instance, immunofluorescent staining and flow microfluorimetry, peptide-dependent class I assembly assays, and inhibition of CTL recognition by peptide competition. Those peptides that bind to the class I molecule are further evaluated for their ability to serve as targets for CTLs derived from infected or immunized individuals, as well as for their capacity to induce primary in vitro or in vivo CTL responses that can give rise to CTL populations capable of reacting with virally infected target cells or tumor cells as potential therapeutic agents.

The MHC class I antigens are encoded by the HLA-A, B, and C loci. HLA-A and B antigens are expressed at the cell surface at approximately equal densities, whereas the expression of HLA-C is significantly lower (perhaps as much as 10-fold lower). Each of these loci have a number of alleles. The peptide binding motifs of the invention are relatively specific for each allelic subtype.

For peptide-based vaccines, the peptides of the present invention preferably comprise a motif recognized by an MHC I molecule having a wide distribution in the human population. Since the MHC alleles occur at different frequencies within different ethnic groups and races, the choice of target MHC allele may depend upon the target population. Table 4 shows the frequency of various alleles at the HLA-A locus products among different races. For instance, the majority of the Caucasoid population can be covered by peptides which bind to four HLA-A allele subtypes, specifically HLA-A2.1, A1, A3.2, and A24.1. Similarly, the majority of the Asian population is encompassed with the addition of peptides binding to a fifth allele HLA-A11.2.

TABLE 4

<u>A Allele/Subtype</u>	<u>N(69)*</u>	<u>A(54)</u>	<u>C(502)</u>
A1	10.1(7)	1.8(1)	27.4(138)
A2.1	11.5(8)	37.0(20)	39.8(199)
A2.2	10.1(7)	0	3.3(17)

A2.3	1.4(1)	5.5(3)	0.8(4)
A2.4	-	-	-
A2.5	-	-	-
A3.1	1.4(1)	0	0.2(0)
A3.2	5.7(4)	5.5(3)	21.5(108)
A11.1	0	5.5(3)	0
A11.2	5.7(4)	31.4(17)	8.7(44)
A11.3	0	3.7(2)	0
A23	4.3(3)	-	3.9(20)
A24	2.9(2)	27.7(15)	15.3(77)
A24.2	-	-	-
A24.3	-	-	-
A25	1.4(1)	-	6.9(35)
A26.1	4.3(3)	9.2(5)	5.9(30)
A26.2	7.2(5)	-	1.0(5)
A26V	-	3.7(2)	-
A28.1	10.1(7)	-	1.6(8)
A28.2	1.4(1)	-	7.5(38)
A29.1	1.4(1)	-	1.4(7)
A29.2	10.1(7)	1.8(1)	5.3(27)
A30.1	8.6(6)	-	4.9(25)
A30.2	1.4(1)	-	0.2(1)
A30.3	7.2(5)	-	3.9(20)
A31	4.3(3)	7.4(4)	6.9(35)
A32	2.8(2)	-	7.1(36)
Aw33.1	8.6(6)	-	2.5(13)
Aw33.2	2.8(2)	16.6(9)	1.2(6)
Aw34.1	1.4(1)	-	-
Aw34.2	14.5(10)	-	0.8(4)
Aw36	5.9(4)	-	-

Table compiled from B. DuPont, Immunobiology of HLA, Vol. I, Histocompatibility Testing 1987, Springer-Verlag, New York 1989.

\* N - negroid; A = Asian; C = Caucasoid. Numbers in parenthesis represent the number of individuals included in the analysis.

5           The nomenclature used to describe peptide compounds follows the conventional practice wherein the amino group is presented to the left (the N-terminus) and the carboxyl group to the right (the C-terminus) of each amino acid residue. In the formulae representing selected specific embodiments of the present invention, the amino- and carboxyl-terminal groups, although not specifically shown, are in the form they  
10       would assume at physiologic pH values, unless otherwise specified. In the amino acid structure formulae, each residue is generally represented by standard three letter or single letter designations. The L-form of an amino acid residue is represented by a capital single letter or a capital first letter of a three-letter symbol, and the D-form for those amino acids is represented by a lower case single letter or a lower case three letter symbol. Glycine  
15       has no asymmetric carbon atom and is simply referred to as "Gly" or G.

          The procedures used to identify peptides of the present invention generally follow the methods disclosed in Falk et al., Nature 351:290 (1991), which is incorporated herein by reference. Briefly, the methods involve large-scale isolation of MHC class I molecules, typically by immunoprecipitation or affinity chromatography, from the  
20       appropriate cell or cell line. Examples of other methods for isolation of the desired MHC molecule equally well known to the artisan include ion exchange chromatography, lectin chromatography, size exclusion, high performance ligand chromatography, and a combination of all of the above techniques.

          A large number of cells with defined MHC molecules, particularly MHC  
25       Class I molecules, are known and readily available. For example, human EBV-transformed B cell lines have been shown to be excellent sources for the preparative isolation of class I and class II MHC molecules. Well-characterized cell lines are available from private and commercial sources, such as American Type Culture Collection ("Catalogue of Cell Lines and Hybridomas," 6th edition (1988) Rockville,  
30       Maryland, U.S.A.); National Institute of General Medical Sciences 1990/1991 Catalog of Cell Lines (NIGMS) Human Genetic Mutant Cell Repository, Camden, NJ; and ASHI Repository, Bingham and Women's Hospital, 75 Francis Street, Boston, MA 02115. Table 5 lists some B cell lines suitable for use as sources for HLA-A alleles. All of these cell lines can be grown in large batches and are therefore useful for large scale production

of MHC molecules. One of skill will recognize that these are merely exemplary cell lines and that many other cell sources can be employed. Similar EBV B cell lines homozygous for HLA-B and HLA-C could serve as sources for HLA-B and HLA-C alleles, respectively.

5

TABLE 5

## HUMAN CELL LINES (HLA-A SOURCES)

HLA-A allele	B cell line
A1	MAT COX (9022) STEINLIN (9087)
A2.1	JY
A3.2	EHM (9080) HO301 (9055)GM3107
A24.1	T3(9107),TISI (9042)
A11	BVR (GM6828A) WT100 (GM8602)WT52 (GM8603)

In the typical case, immunoprecipitation is used to isolate the desired allele. A number of protocols can be used, depending upon the specificity of the antibodies used. For example, allele-specific mAb reagents can be used for the affinity purification of the HLA-A, HLA-B, and HLA-C molecules. Several mAb reagents for the isolation of HLA-A molecules are available (Table 6). Thus, for each of the targeted HLA-A alleles, reagents are available that may be used for the direct isolation of the HLA-A molecules. Affinity columns prepared with these mAbs using standard techniques are successfully used to purify the respective HLA-A allele products.

15

In addition to allele-specific mAbs, broadly reactive anti-HLA-A, B, C mAbs, such as W6/32 and B9.12.1, and one anti-HLA-B, C mAb, B1.23.2, could be used in alternative affinity purification protocols as described in the example section below.

TABLE 6

## ANTIBODY REAGENTS

anti-HLA	Name	
HLA-A1	12/18	
HLA-A3	GAPA3	(ATCC, HB122)
HLA-11,24.1	A11.1M	(ATCC, HB164)
HLA-A,B,C	W6/32	(ATCC, HB95)
monomorphic	B9.12.1	(INSERM-CNRS)
HLA-B,C	B.1.23.2	(INSERM-CNRS)
monomorphic		

5                   The peptides bound to the peptide binding groove of the isolated MHC molecules are eluted typically using acid treatment. Peptides can also be dissociated from class I molecules by a variety of standard denaturing means, such as heat, pH, detergents, salts, chaotropic agents, or a combination thereof.

10                   Peptide fractions are further separated from the MHC molecules by reversed-phase high performance liquid chromatography (HPLC) and sequenced. Peptides can be separated by a variety of other standard means well known to the artisan, including filtration, ultrafiltration, electrophoresis, size chromatography, precipitation with specific antibodies, ion exchange chromatography, isoelectrofocusing, and the like.

15                   Sequencing of the isolated peptides can be performed according to standard techniques such as Edman degradation (Hunkapiller, M.W., et al., Methods Enzymol. 91, 399 [1983]). Other methods suitable for sequencing include mass spectrometry sequencing of individual peptides as previously described (Hunt, et al., Science 225:1261 (1992), which is incorporated herein by reference). Amino acid sequencing of bulk heterogenous peptides (e.g., pooled HPLC fractions) from different class I molecules typically reveals a characteristic sequence motif for each class I allele.

20                   Definition of motifs specific for different class I alleles allows the identification of potential peptide epitopes from an antigenic protein whose amino acid sequence is known. Typically, identification of potential peptide epitopes is initially



carried out using a computer to scan the amino acid sequence of a desired antigen for the presence of motifs. The epitopic sequences are then synthesized. The capacity to bind MHC Class molecules is measured in a variety of different ways. One means is a Class I molecule binding assay as described in the related applications, noted above. Other  
 5 alternatives described in the literature include inhibition of antigen presentation (Sette, et al., J. Immunol. 141:3893 (1991), in vitro assembly assays (Townsend, et al., Cell 62:285 (1990), and FACS based assays using mutated cells, such as RMA.S (Melief, et al., Eur. J. Immunol. 21:2963 (1991)).

Next, peptides that test positive in the MHC class I binding assay are  
 10 assayed for the ability of the peptides to induce specific CTL responses in vitro. For instance, Antigen-presenting cells that have been incubated with a peptide can be assayed for the ability to induce CTL responses in responder cell populations. Antigen-presenting cells can be normal cells such as peripheral blood mononuclear cells or dendritic cells (Inaba, et al., J. Exp. Med. 166:182 (1987); Boog, Eur. J. Immunol. 18:219 [1988]).

Alternatively, mutant mammalian cell lines that are deficient in their  
 15 ability to load class I molecules with internally processed peptides, such as the mouse cell lines RMA-S (Kärre, et al., Nature, 319:675 (1986); Ljunggren, et al., Eur. J. Immunol. 21:2963-2970 (1991)), and the human somatic T cell hybrid, T-2 (Cerundolo, et al., Nature 345:449-452 (1990)) and which have been transfected with the appropriate human  
 20 class I genes are conveniently used, when peptide is added to them, to test for the capacity of the peptide to induce in vitro primary CTL responses. Other eukaryotic cell lines which could be used include various insect cell lines such as mosquito larvae (ATCC cell lines CCL 125, 126, 1660, 1591, 6585, 6586), silkworm (ATTC CRL 8851), armyworm (ATCC CRL 1711), moth (ATCC CCL 80) and Drosophila cell lines such as a Schneider  
 25 cell line (see Schneider J. Embryol. Exp. Morphol. 27:353-365 [1927]).

Peripheral blood lymphocytes are conveniently isolated following simple venipuncture or leukapheresis of normal donors or patients and used as the responder cell sources of CTL precursors. In one embodiment, the appropriate antigen-presenting cells are incubated with 10-100  $\mu$ M of peptide in serum-free media for 4 hours under  
 30 appropriate culture conditions. The peptide-loaded antigen-presenting cells are then incubated with the responder cell populations in vitro for 7 to 10 days under optimized culture conditions. Positive CTL activation can be determined by assaying the cultures for the presence of CTLs that kill radiolabeled target cells, both specific peptide-pulsed

targets as well as target cells expressing endogenously processed form of the relevant virus or tumor antigen from which the peptide sequence was derived.

Specificity and MHC restriction of the CTL is determined by testing against different peptide target cells expressing appropriate or inappropriate human MHC class I. The peptides that test positive in the MHC binding assays and give rise to specific CTL responses are referred to herein as immunogenic peptides.

The immunogenic peptides can be prepared synthetically, or by recombinant DNA technology or from natural sources such as whole viruses or tumors. Although the peptide will preferably be substantially free of other naturally occurring host cell proteins and fragments thereof, in some embodiments the peptides can be synthetically conjugated to native fragments or particles.

The polypeptides or peptides can be a variety of lengths, either in their neutral (uncharged) forms or in forms which are salts, and either free of modifications such as glycosylation, side chain oxidation, or phosphorylation or containing these modifications, subject to the condition that the modification not destroy the biological activity of the polypeptides as herein described.

Desirably, the peptide will be as small as possible while still maintaining substantially all of the biological activity of the large peptide. When possible, it may be desirable to optimize peptides of the invention to a length of 9 or 10 amino acid residues, commensurate in size with endogenously processed viral peptides or tumor cell peptides that are bound to MHC class I molecules on the cell surface.

Peptides having the desired activity may be modified as necessary to provide certain desired attributes, e.g., improved pharmacological characteristics, while increasing or at least retaining substantially all of the biological activity of the unmodified peptide to bind the desired MHC molecule and activate the appropriate T cell. For instance, the peptides may be subject to various changes, such as substitutions, either conservative or non-conservative, where such changes might provide for certain advantages in their use, such as improved MHC binding. By conservative substitutions is meant replacing an amino acid residue with another which is biologically and/or chemically similar, e.g., one hydrophobic residue for another, or one polar residue for another. The substitutions include combinations such as Gly, Ala; Val, Ile, Leu, Met; Asp, Glu; Asn, Gln; Ser, Thr; Lys, Arg; and Phe, Tyr. The effect of single amino acid substitutions may also be probed using D-amino acids. Such modifications may be made

using well known peptide synthesis procedures, as described in e.g., Merrifield, Science 232:341-347 (1986), Barany and Merrifield, The Peptides, Gross and Meienhofer, eds. (N.Y., Academic Press), pp. 1-284 (1979); and Stewart and Young, Solid Phase Peptide Synthesis, (Rockford, Ill., Pierce), 2d Ed. (1984), incorporated by reference herein.

5                   The peptides can also be modified by extending or decreasing the compound's amino acid sequence, e.g., by the addition or deletion of amino acids. The peptides or analogs of the invention can also be modified by altering the order or composition of certain residues, it being readily appreciated that certain amino acid residues essential for biological activity, e.g., those at critical contact sites or conserved  
10 residues, may generally not be altered without an adverse effect on biological activity. The non-critical amino acids need not be limited to those naturally occurring in proteins, such as L- $\alpha$ -amino acids, or their D-isomers, but may include non-natural amino acids as well, such as  $\beta$ - $\gamma$ - $\delta$ -amino acids, as well as many derivatives of L- $\alpha$ -amino acids.

Typically, a series of peptides with single amino acid substitutions are  
15 employed to determine the effect of electrostatic charge, hydrophobicity, etc. on binding. For instance, a series of positively charged (e.g., Lys or Arg) or negatively charged (e.g., Glu) amino acid substitutions are made along the length of the peptide revealing different patterns of sensitivity towards various MHC molecules and T cell receptors. In addition, multiple substitutions using small, relatively neutral moieties such as Ala, Gly, Pro, or  
20 similar residues may be employed. The substitutions may be homo-oligomers or hetero-oligomers. The number and types of residues which are substituted or added depend on the spacing necessary between essential contact points and certain functional attributes which are sought (e.g., hydrophobicity versus hydrophilicity). Increased binding affinity for an MHC molecule or T cell receptor may also be achieved by such substitutions,  
25 compared to the affinity of the parent peptide. In any event, such substitutions should employ amino acid residues or other molecular fragments chosen to avoid, for example, steric and charge interference which might disrupt binding.

Amino acid substitutions are typically of single residues. Substitutions, deletions, insertions or any combination thereof may be combined to arrive at a final  
30 peptide. Substitutional variants are those in which at least one residue of a peptide has been removed and a different residue inserted in its place. Such substitutions generally are made in accordance with the following Table 2 when it is desired to finely modulate the characteristics of the peptide.

TABLE 2

<u>Original Residue</u>	<u>Exemplary Substitution</u>
Ala	Ser
Arg	Lys, His
Asn	Gln
Asp	Glu
Cys	Ser
Glu	Asp
Gly	Pro
His	Lys; Arg
Ile	Leu; Val
Leu	Ile; Val
Lys	Arg; His
Met	Leu; Ile
Phe	Tyr; Trp
Ser	Thr
Thr	Ser
Trp	Tyr; Phe
Tyr	Trp; Phe
Val	Ile; Leu
Pro	Gly

Substantial changes in function (e.g., affinity for MHC molecules or T cell  
5 receptors) are made by selecting substitutions that are less conservative than those in  
Table 2, i.e., selecting residues that differ more significantly in their effect on maintaining  
(a) the structure of the peptide backbone in the area of the substitution, for example as a  
sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the  
target site or (c) the bulk of the side chain. The substitutions which in general are  
10 expected to produce the greatest changes in peptide properties will be those in which (a)  
hydrophilic residue, e.g. seryl, is substituted for (or by) a hydrophobic residue, e.g. leucyl,  
isoleucyl, phenylalanyl, valyl or alanyl; (b) a residue having an electropositive side chain,  
e.g., lysyl, arginyl, or histidyl, is substituted for (or by) an electronegative residue, e.g.

glutamyl or aspartyl; or (c) a residue having a bulky side chain, e.g. phenylalanine, is substituted for (or by) one not having a side chain, e.g., glycine.

The peptides may also comprise isosteres of two or more residues in the immunogenic peptide. An isostere as defined here is a sequence of two or more residues that can be substituted for a second sequence because the steric conformation of the first sequence fits a binding site specific for the second sequence. The term specifically includes peptide backbone modifications well known to those skilled in the art. Such modifications include modifications of the amide nitrogen, the  $\alpha$ -carbon, amide carbonyl, complete replacement of the amide bond, extensions, deletions or backbone crosslinks.

See, generally, Spatola, *Chemistry and Biochemistry of Amino Acids, peptides and Proteins*, Vol. VII (Weinstein ed., 1983).

Modifications of peptides with various amino acid mimetics or unnatural amino acids are particularly useful in increasing the stability of the peptide in vivo. Stability can be assayed in a number of ways. For instance, peptidases and various biological media, such as human plasma and serum, have been used to test stability. See, e.g., Verhoef et al., Eur. J. Drug Metab. Pharmacokin. 11:291-302 (1986). Half life of the peptides of the present invention is conveniently determined using a 25% human serum (v/v) assay. The protocol is generally as follows. Pooled human serum (Type AB, non-heat inactivated) is delipidated by centrifugation before use. The serum is then diluted to 25% with RPMI tissue culture media and used to test peptide stability. At predetermined time intervals a small amount of reaction solution is removed and added to either 6% aqueous trichloroacetic acid or ethanol. The cloudy reaction sample is cooled (4°C) for 15 minutes and then spun to pellet the precipitated serum proteins. The presence of the peptides is then determined by reversed-phase HPLC using stability-specific chromatography conditions.

The peptides of the present invention or analogs thereof which have CTL stimulating activity may be modified to provide desired attributes other than improved serum half life. For instance, the ability of the peptides to induce CTL activity can be enhanced by linkage to a sequence which contains at least one epitope that is capable of inducing a T helper cell response. Particularly preferred immunogenic peptides/T helper conjugates are linked by a spacer molecule. The spacer is typically comprised of relatively small, neutral molecules, such as amino acids or amino acid mimetics, which are substantially uncharged under physiological conditions. The spacers are typically

selected from, e.g., Ala, Gly, or other neutral spacers of nonpolar amino acids or neutral polar amino acids. It will be understood that the optionally present spacer need not be comprised of the same residues and thus may be a hetero- or homo-oligomer. When present, the spacer will usually be at least one or two residues, more usually three to six residues. Alternatively, the CTL peptide may be linked to the T helper peptide without a spacer.

The immunogenic peptide may be linked to the T helper peptide either directly or via a spacer either at the amino or carboxy terminus of the CTL peptide. The amino terminus of either the immunogenic peptide or the T helper peptide may be acylated. Exemplary T helper peptides include tetanus toxoid 830-843, influenza 307-319, malaria circumsporozoite 382-398 and 378-389.

In some embodiments it may be desirable to include in the pharmaceutical compositions of the invention at least one component which primes CTL. Lipids have been identified as agents capable of priming CTL in vivo against viral antigens. For example, palmitic acid residues can be attached to the alpha and epsilon amino groups of a Lys residue and then linked, e.g., via one or more linking residues such as Gly, Gly-Gly-, Ser, Ser-Ser, or the like, to an immunogenic peptide. The lipidated peptide can then be injected directly in a micellar form, incorporated into a liposome or emulsified in an adjuvant, e.g., incomplete Freund's adjuvant. In a preferred embodiment a particularly effective immunogen comprises palmitic acid attached to alpha and epsilon amino groups of Lys, which is attached via linkage, e.g., Ser-Ser, to the amino terminus of the immunogenic peptide.

As another example of lipid priming of CTL responses, *E. coli* lipoproteins, such as tripalmitoyl-S-glycerylcysteinylserine (P3CSS) can be used to prime virus specific CTL when covalently attached to an appropriate peptide. See, Deres et al., Nature 342:561-564 (1989), incorporated herein by reference. Peptides of the invention can be coupled to P<sup>3</sup>CSS, for example, and the lipopeptide administered to an individual to specifically prime a CTL response to the target antigen. Further, as the induction of neutralizing antibodies can also be primed with P<sup>3</sup>CSS conjugated to a peptide which displays an appropriate epitope, the two compositions can be combined to more effectively elicit both humoral and cell-mediated responses to infection.

In addition, additional amino acids can be added to the termini of a peptide to provide for ease of linking peptides one to another, for coupling to a carrier support, or

larger peptide, for modifying the physical or chemical properties of the peptide or oligopeptide, or the like. Amino acids such as tyrosine, cysteine, lysine, glutamic or aspartic acid, or the like, can be introduced at the C- or N-terminus of the peptide or oligopeptide. Modification at the C terminus in some cases may alter binding characteristics of the peptide. In addition, the peptide or oligopeptide sequences can differ from the natural sequence by being modified by terminal-NH<sub>2</sub>-acylation, e.g., by alkanoyl (C<sub>1</sub>-C<sub>20</sub>) or thioglycolyl acetylation, terminal-carboxyl amidation, e.g., ammonia, methylamine, etc. In some instances these modifications may provide sites for linking to a support or other molecule.

The peptides of the invention can be prepared in a wide variety of ways. Because of their relatively short size, the peptides can be synthesized in solution or on a solid support in accordance with conventional techniques. Various automatic synthesizers are commercially available and can be used in accordance with known protocols. See, for example, Stewart and Young, Solid Phase Peptide Synthesis, 2d. ed., Pierce Chemical Co. (1984), supra.

Alternatively, recombinant DNA technology may be employed wherein a nucleotide sequence which encodes an immunogenic peptide of interest is inserted into an expression vector, transformed or transfected into an appropriate host cell and cultivated under conditions suitable for expression. These procedures are generally known in the art, as described generally in Sambrook et al., Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Cold Spring Harbor, New York (1982), which is incorporated herein by reference. Thus, fusion proteins which comprise one or more peptide sequences of the invention can be used to present the appropriate T cell epitope.

As the coding sequence for peptides of the length contemplated herein can be synthesized by chemical techniques, for example, the phosphotriester method of Matteucci et al., J. Am. Chem. Soc. 103:3185 (1981), modification can be made simply by substituting the appropriate base(s) for those encoding the native peptide sequence. The coding sequence can then be provided with appropriate linkers and ligated into expression vectors commonly available in the art, and the vectors used to transform suitable hosts to produce the desired fusion protein. A number of such vectors and suitable host systems are now available. For expression of the fusion proteins, the coding sequence will be provided with operably linked start and stop codons, promoter and terminator regions and usually a replication system to provide an expression vector for

expression in the desired cellular host. For example, promoter sequences compatible with bacterial hosts are provided in plasmids containing convenient restriction sites for insertion of the desired coding sequence. The resulting expression vectors are transformed into suitable bacterial hosts. Of course, yeast or mammalian cell hosts may also be used, employing suitable vectors and control sequences.

The peptides of the present invention and pharmaceutical and vaccine compositions thereof are useful for administration to mammals, particularly humans, to treat and/or prevent viral infection and cancer. Examples of diseases which can be treated using the immunogenic peptides of the invention include prostate cancer, hepatitis B, hepatitis C, AIDS, renal carcinoma, cervical carcinoma, lymphoma, CMV and condyloma acuminatum.

For pharmaceutical compositions, the immunogenic peptides of the invention are administered to an individual already suffering from cancer or infected with the virus of interest. Those in the incubation phase or the acute phase of infection can be treated with the immunogenic peptides separately or in conjunction with other treatments, as appropriate. In therapeutic applications, compositions are administered to a patient in an amount sufficient to elicit an effective CTL response to the virus or tumor antigen and to cure or at least partially arrest symptoms and/or complications. An amount adequate to accomplish this is defined as "therapeutically effective dose." Amounts effective for this use will depend on, e.g., the peptide composition, the manner of administration, the stage and severity of the disease being treated, the weight and general state of health of the patient, and the judgment of the prescribing physician, but generally range for the initial immunization (that is for therapeutic or prophylactic administration) from about 1.0  $\mu\text{g}$  to about 5000  $\mu\text{g}$  of peptide for a 70 kg patient, followed by boosting dosages of from about 1.0  $\mu\text{g}$  to about 1000  $\mu\text{g}$  of peptide pursuant to a boosting regimen over weeks to months depending upon the patient's response and condition by measuring specific CTL activity in the patient's blood. It must be kept in mind that the peptides and compositions of the present invention may generally be employed in serious disease states, that is, life-threatening or potentially life threatening situations. In such cases, in view of the minimization of extraneous substances and the relative nontoxic nature of the peptides, it is possible and may be felt desirable by the treating physician to administer substantial excesses of these peptide compositions.



For therapeutic use, administration should begin at the first sign of viral infection or the detection or surgical removal of tumors or shortly after diagnosis in the case of acute infection. This is followed by boosting doses until at least symptoms are substantially abated and for a period thereafter. In chronic infection, loading doses followed by boosting doses may be required.

Treatment of an infected individual with the compositions of the invention may hasten resolution of the infection in acutely infected individuals. For those individuals susceptible (or predisposed) to developing chronic infection the compositions are particularly useful in methods for preventing the evolution from acute to chronic infection. Where the susceptible individuals are identified prior to or during infection, for instance, as described herein, the composition can be targeted to them, minimizing need for administration to a larger population.

The peptide compositions can also be used for the treatment of chronic infection and to stimulate the immune system to eliminate virus-infected cells in carriers. It is important to provide an amount of immuno-potentiating peptide in a formulation and mode of administration sufficient to effectively stimulate a cytotoxic T cell response. Thus, for treatment of chronic infection, a representative dose is in the range of about 1.0 µg to about 5000 µg, preferably about 5 µg to 1000 µg for a 70 kg patient per dose. Immunizing doses followed by boosting doses at established intervals, e.g., from one to four weeks, may be required, possibly for a prolonged period of time to effectively immunize an individual. In the case of chronic infection, administration should continue until at least clinical symptoms or laboratory tests indicate that the viral infection has been eliminated or substantially abated and for a period thereafter.

The pharmaceutical compositions for therapeutic treatment are intended for parenteral, topical, oral or local administration. Preferably, the pharmaceutical compositions are administered parenterally, e.g., intravenously, subcutaneously, intradermally, or intramuscularly. Thus, the invention provides compositions for parenteral administration which comprise a solution of the immunogenic peptides dissolved or suspended in an acceptable carrier, preferably an aqueous carrier. A variety of aqueous carriers may be used, e.g., water, buffered water, 0.8% saline, 0.3% glycine, hyaluronic acid and the like. These compositions may be sterilized by conventional, well known sterilization techniques, or may be sterile filtered. The resulting aqueous solutions may be packaged for use as is, or lyophilized, the lyophilized preparation being combined

with a sterile solution prior to administration. The compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions, such as pH adjusting and buffering agents, tonicity adjusting agents, wetting agents and the like, for example, sodium acetate, sodium lactate, sodium chloride, potassium chloride, calcium chloride, sorbitan monolaurate, triethanolamine oleate, etc.

The concentration of CTL stimulatory peptides of the invention in the pharmaceutical formulations can vary widely, i.e., from less than about 0.1%, usually at or at least about 2% to as much as 20% to 50% or more by weight, and will be selected primarily by fluid volumes, viscosities, etc., in accordance with the particular mode of administration selected.

The peptides of the invention may also be administered via liposomes, which serve to target the peptides to a particular tissue, such as lymphoid tissue, or targeted selectively to infected cells, as well as increase the half-life of the peptide composition. Liposomes include emulsions, foams, micelles, insoluble monolayers, liquid crystals, phospholipid dispersions, lamellar layers and the like. In these preparations the peptide to be delivered is incorporated as part of a liposome, alone or in conjunction with a molecule which binds to, e.g., a receptor prevalent among lymphoid cells, such as monoclonal antibodies which bind to the CD45 antigen, or with other therapeutic or immunogenic compositions. Thus, liposomes either filled or decorated with a desired peptide of the invention can be directed to the site of lymphoid cells, where the liposomes then deliver the selected therapeutic/immunogenic peptide compositions. Liposomes for use in the invention are formed from standard vesicle-forming lipids, which generally include neutral and negatively charged phospholipids and a sterol, such as cholesterol. The selection of lipids is generally guided by consideration of, e.g., liposome size, acid lability and stability of the liposomes in the blood stream. A variety of methods are available for preparing liposomes, as described in, e.g., Szoka et al., Ann. Rev. Biophys. Bioeng. 9:467 (1980), U.S. Patent Nos. 4,235,871, 4,501,728, 4,837,028, and 5,019,369, incorporated herein by reference.

For targeting to the immune cells, a ligand to be incorporated into the liposome can include, e.g., antibodies or fragments thereof specific for cell surface determinants of the desired immune system cells. A liposome suspension containing a peptide may be administered intravenously, locally, topically, etc. in a dose which varies

according to, inter alia, the manner of administration, the peptide being delivered, and the stage of the disease being treated.

For solid compositions, conventional nontoxic solid carriers may be used which include, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharin, talcum, cellulose, glucose, sucrose, magnesium carbonate, and the like. For oral administration, a pharmaceutically acceptable nontoxic composition is formed by incorporating any of the normally employed excipients, such as those carriers previously listed, and generally 10-95% of active ingredient, that is, one or more peptides of the invention, and more preferably at a concentration of 25%-75%.

For aerosol administration, the immunogenic peptides are preferably supplied in finely divided form along with a surfactant and propellant. Typical percentages of peptides are 0.01%-20% by weight, preferably 1%-10%. The surfactant must, of course, be nontoxic, and preferably soluble in the propellant. Representative of such agents are the esters or partial esters of fatty acids containing from 6 to 22 carbon atoms, such as caproic, octanoic, lauric, palmitic, stearic, linoleic, linolenic, olesteric and oleic acids with an aliphatic polyhydric alcohol or its cyclic anhydride. Mixed esters, such as mixed or natural glycerides may be employed. The surfactant may constitute 0.1%-20% by weight of the composition, preferably 0.25-5%. The balance of the composition is ordinarily propellant. A carrier can also be included, as desired, as with, e.g., lecithin for intranasal delivery.

In another aspect the present invention is directed to vaccines which contain as an active ingredient an immunogenically effective amount of an immunogenic peptide as described herein. The peptide(s) may be introduced into a host, including humans, linked to its own carrier or as a homopolymer or heteropolymer of active peptide units. Such a polymer has the advantage of increased immunological reaction and, where different peptides are used to make up the polymer, the additional ability to induce antibodies and/or CTLs that react with different antigenic determinants of the virus or tumor cells. Useful carriers are well known in the art, and include, e.g., thyroglobulin, albumins such as human serum albumin, tetanus toxoid, polyamino acids such as poly(lysine:glutamic acid), influenza, hepatitis B virus core protein, hepatitis B virus recombinant vaccine and the like. The vaccines can also contain a physiologically tolerable (acceptable) diluent such as water, phosphate buffered saline, or saline, and further typically include an adjuvant. Adjuvants such as incomplete Freund's adjuvant,

aluminum phosphate, aluminum hydroxide, or alum are materials well known in the art. And, as mentioned above, CTL responses can be primed by conjugating peptides of the invention to lipids, such as P<sub>3</sub>CSS. Upon immunization with a peptide composition as described herein, via injection, aerosol, oral, transdermal or other route, the immune system of the host responds to the vaccine by producing large amounts of CTLs specific for the desired antigen, and the host becomes at least partially immune to later infection, or resistant to developing chronic infection.

Vaccine compositions containing the peptides of the invention are administered to a patient susceptible to or otherwise at risk of viral infection or cancer to elicit an immune response against the antigen and thus enhance the patient's own immune response capabilities. Such an amount is defined to be an "immunogenically effective dose." In this use, the precise amounts again depend on the patient's state of health and weight, the mode of administration, the nature of the formulation, etc., but generally range from about 1.0 µg to about 5000 µg per 70 kilogram patient, more commonly from about 10 µg to about 500 µg mg per 70 kg of body weight.

In some instances it may be desirable to combine the peptide vaccines of the invention with vaccines which induce neutralizing antibody responses to the virus of interest, particularly to viral envelope antigens.

For therapeutic or immunization purposes, nucleic acids encoding one or more of the peptides of the invention can also be administered to the patient. A number of methods are conveniently used to deliver the nucleic acids to the patient. For instance, the nucleic acid can be delivered directly, as "naked DNA". This approach is described, for instance, in Wolff et. al., *Science* 247:1465-1468 (1990) as well as U.S. Patent Nos. 5,580,859 and 5,589,466. The nucleic acids can also be administered using ballistic delivery as described, for instance, in U.S. Patent No. 5,204,253. Particles comprised solely of DNA can be administered. Alternatively, DNA can be adhered to particles, such as gold particles. The nucleic acids can also be delivered complexed to cationic compounds, such as cationic lipids. Lipid-mediated gene delivery methods are described, for instance, in WO 96/18372; WO 93/24640; Mannino and Gould-Fogerite (1988) *BioTechniques* 6(7):682-691; Rose U.S. Pat No. 5,279,833; WO 91/06309; and Felgner et al. (1987) *Proc. Natl. Acad. Sci. USA* 84: 7413-7414. The peptides of the invention can also be expressed by attenuated viral hosts, such as vaccinia or fowlpox. This approach involves the use of vaccinia virus as a vector to express nucleotide sequences that encode

the peptides of the invention. Upon introduction into an acutely or chronically infected host or into a noninfected host, the recombinant vaccinia virus expresses the immunogenic peptide, and thereby elicits a host CTL response. Vaccinia vectors and methods useful in immunization protocols are described in, e.g., U.S. Patent No.

5 4,722,848, incorporated herein by reference. Another vector is BCG (Bacille Calmette Guerin). BCG vectors are described in Stover et al. (Nature 351:456-460 (1991)) which is incorporated herein by reference. A wide variety of other vectors useful for therapeutic administration or immunization of the peptides of the invention, e.g., *Salmonella typhi* vectors and the like, will be apparent to those skilled in the art from the description  
10 herein.

A preferred means of administering nucleic acids encoding the peptides of the invention uses minigene constructs encoding multiple epitopes of the invention. To create a DNA sequence encoding the selected CTL epitopes (minigene) for expression in human cells, the amino acid sequences of the epitopes are reverse translated. A human  
15 codon usage table is used to guide the codon choice for each amino acid. These epitope-encoding DNA sequences are directly adjoined, creating a continuous polypeptide sequence. To optimize expression and/or immunogenicity, additional elements can be incorporated into the minigene design. Examples of amino acid sequence that could be reverse translated and included in the minigene sequence include: helper T lymphocyte  
20 epitopes, a leader (signal) sequence, and an endoplasmic reticulum retention signal. In addition, MHC presentation of CTL epitopes may be improved by including synthetic (e.g. poly-alanine) or naturally-occurring flanking sequences adjacent to the CTL epitopes.

The minigene sequence is converted to DNA by assembling  
25 oligonucleotides that encode the plus and minus strands of the minigene. Overlapping oligonucleotides (30-100 bases long) are synthesized, phosphorylated, purified and annealed under appropriate conditions using well known techniques. The ends of the oligonucleotides are joined using T4 DNA ligase. This synthetic minigene, encoding the CTL epitope polypeptide, can then be cloned into a desired expression vector.

30 Standard regulatory sequences well known to those of skill in the art are included in the vector to ensure expression in the target cells. Several vector elements are required: a promoter with a down-stream cloning site for minigene insertion; a polyadenylation signal for efficient transcription termination; an *E. coli* origin of

replication; and an *E. coli* selectable marker (e.g. ampicillin or kanamycin resistance). Numerous promoters can be used for this purpose, e.g., the human cytomegalovirus (hCMV) promoter. *See*, U.S. Patent Nos. 5,580,859 and 5,589,466 for other suitable promoter sequences.

5 Additional vector modifications may be desired to optimize minigene expression and immunogenicity. In some cases, introns are required for efficient gene expression, and one or more synthetic or naturally-occurring introns could be incorporated into the transcribed region of the minigene. The inclusion of mRNA stabilization sequences can also be considered for increasing minigene expression. It has recently been proposed that immunostimulatory sequences (ISSs or CpGs) play a role in the immunogenicity of DNA vaccines. These sequences could be included in the vector, outside the minigene coding sequence, if found to enhance immunogenicity.

In some embodiments, a bioisotonic expression vector, to allow production of the minigene-encoded epitopes and a second protein included to enhance or decrease immunogenicity can be used. Examples of proteins or polypeptides that could beneficially enhance the immune response if co-expressed include cytokines (e.g., IL2, IL12, GM-CSF), cytokine-inducing molecules (e.g. LeIF) or costimulatory molecules. Helper (HTL) epitopes could be joined to intracellular targeting signals and expressed separately from the CTL epitopes. This would allow direction of the HTL epitopes to a cell compartment different than the CTL epitopes. If required, this could facilitate more efficient entry of HTL epitopes into the MHC class II pathway, thereby improving CTL induction. In contrast to CTL induction, specifically decreasing the immune response by co-expression of immunosuppressive molecules (e.g. TGF- $\beta$ ) may be beneficial in certain diseases.

25 Once an expression vector is selected, the minigene is cloned into the polylinker region downstream of the promoter. This plasmid is transformed into an appropriate *E. coli* strain, and DNA is prepared using standard techniques. The orientation and DNA sequence of the minigene, as well as all other elements included in the vector, are confirmed using restriction mapping and DNA sequence analysis.

30 Bacterial cells harboring the correct plasmid can be stored as a master cell bank and a working cell bank.

Therapeutic quantities of plasmid DNA are produced by fermentation in *E. coli*, followed by purification. Aliquots from the working cell bank are used to inoculate

fermentation medium (such as Terrific Broth), and grown to saturation in shaker flasks or a bioreactor according to well known techniques. Plasmid DNA can be purified using standard bioseparation technologies such as solid phase anion-exchange resins supplied by Quiagen. If required, supercoiled DNA can be isolated from the open circular and  
5 linear forms using gel electrophoresis or other methods.

Purified plasmid DNA can be prepared for injection using a variety of formulations. The simplest of these is reconstitution of lyophilized DNA in sterile phosphate-buffer saline (PBS). A variety of methods have been described, and new techniques may become available. As noted above, nucleic acids are conveniently  
10 formulated with cationic lipids. In addition, glycolipids, fusogenic liposomes, peptides and compounds referred to collectively as protective, interactive, non-condensing (PINC) could also be complexed to purified plasmid DNA to influence variables such as stability, intramuscular dispersion, or trafficking to specific organs or cell types.

Target cell sensitization can be used as a functional assay for expression  
15 and MHC class I presentation of minigene-encoded CTL epitopes. The plasmid DNA is introduced into a mammalian cell line that is suitable as a target for standard CTL chromium release assays. The transfection method used will be dependent on the final formulation. Electroporation can be used for "naked" DNA, whereas cationic lipids allow direct in vitro transfection. A plasmid expressing green fluorescent protein (GFP) can be  
20 co-transfected to allow enrichment of transfected cells using fluorescence activated cell sorting (FACS). These cells are then chromium-51 labeled and used as target cells for epitope-specific CTL lines. Cytolysis, detected by <sup>51</sup>Cr release, indicates production of MHC presentation of minigene-encoded CTL epitopes.

*In vivo* immunogenicity is a second approach for functional testing of  
25 minigene DNA formulations. Transgenic mice expressing appropriate human MHC molecules are immunized with the DNA product. The dose and route of administration are formulation dependent (e.g. IM for DNA in PBS, IP for lipid-complexed DNA). Twenty-one days after immunization, splenocytes are harvested and restimulated for 1 week in the presence of peptides encoding each epitope being tested. These effector cells  
30 (CTLs) are assayed for cytolysis of peptide-loaded, chromium-51 labeled target cells using standard techniques. Lysis of target cells sensitized by MHC loading of peptides corresponding to minigene-encoded epitopes demonstrates DNA vaccine function for *in vivo* induction of CTLs.

Antigenic peptides may be used to elicit CTL *ex vivo*, as well. The resulting CTL, can be used to treat chronic infections (viral or bacterial) or tumors in patients that do not respond to other conventional forms of therapy, or will not respond to a peptide vaccine approach of therapy. *Ex vivo* CTL responses to a particular pathogen (infectious agent or tumor antigen) are induced by incubating in tissue culture the patient's CTL precursor cells (CTLp) together with a source of antigen-presenting cells (APC) and the appropriate immunogenic peptide. After an appropriate incubation time (typically 1-4 weeks), in which the CTLp are activated and mature and expand into effector CTL, the cells are infused back into the patient, where they will destroy their specific target cell (an infected cell or a tumor cell). In order to optimize the *in vitro* conditions for the generation of specific cytotoxic T cells, the culture of stimulator cells is maintained in an appropriate serum-free medium.

Prior to incubation of the stimulator cells with the cells to be activated, e.g., precursor CD8<sup>+</sup> cells, an amount of antigenic peptide is added to the stimulator cell culture, of sufficient quantity to become loaded onto the human Class I molecules to be expressed on the surface of the stimulator cells. In the present invention, a sufficient amount of peptide is an amount that will allow about 200, and preferably 200 or more, human Class I MHC molecules loaded with peptide to be expressed on the surface of each stimulator cell. Preferably, the stimulator cells are incubated with >20µg/ml peptide.

Resting or precursor CD8<sup>+</sup> cells are then incubated in culture with the appropriate stimulator cells for a time period sufficient to activate the CD8<sup>+</sup> cells. Preferably, the CD8<sup>+</sup> cells are activated in an antigen-specific manner. The ratio of resting or precursor CD8<sup>+</sup> (effector) cells to stimulator cells may vary from individual to individual and may further depend upon variables such as the amenability of an individual's lymphocytes to culturing conditions and the nature and severity of the disease condition or other condition for which the within-described treatment modality is used. Preferably, however, the lymphocyte:stimulator cell ratio is in the range of about 30:1 to 300:1. The effector/stimulator culture may be maintained for as long a time as is necessary to stimulate a therapeutically useable or effective number of CD8<sup>+</sup> cells.

The induction of CTL *in vitro* requires the specific recognition of peptides that are bound to allele specific MHC class I molecules on APC. The number of specific MHC/peptide complexes per APC is crucial for the stimulation of CTL, particularly in primary immune responses. While small amounts of peptide/MHC complexes per cell



are sufficient to render a cell susceptible to lysis by CTL, or to stimulate a secondary CTL response, the successful activation of a CTL precursor (pCTL) during primary response requires a significantly higher number of MHC/peptide complexes. Peptide loading of empty major histocompatibility complex molecules on cells allows the induction of primary cytotoxic T lymphocyte responses. Peptide loading of empty major histocompatibility complex molecules on cells enables the induction of primary cytotoxic T lymphocyte responses.

Since mutant cell lines do not exist for every human MHC allele, it is advantageous to use a technique to remove endogenous MHC-associated peptides from the surface of APC, followed by loading the resulting empty MHC molecules with the immunogenic peptides of interest. The use of non-transformed (non-tumorigenic), non-infected cells, and preferably, autologous cells of patients as APC is desirable for the design of CTL induction protocols directed towards development of *ex vivo* CTL therapies. This application discloses methods for stripping the endogenous MHC-associated peptides from the surface of APC followed by the loading of desired peptides.

A stable MHC class I molecule is a trimeric complex formed of the following elements: 1) a peptide usually of 8 - 10 residues, 2) a transmembrane heavy polymorphic protein chain which bears the peptide-binding site in its  $\alpha 1$  and  $\alpha 2$  domains, and 3) a non-covalently associated non-polymorphic light chain,  $\beta_2$  microglobulin. Removing the bound peptides and/or dissociating the  $\beta_2$  microglobulin from the complex renders the MHC class I molecules nonfunctional and unstable, resulting in rapid degradation. All MHC class I molecules isolated from PBMCs have endogenous peptides bound to them. Therefore, the first step is to remove all endogenous peptides bound to MHC class I molecules on the APC without causing their degradation before exogenous peptides can be added to them.

Two possible ways to free up MHC class I molecules of bound peptides include lowering the culture temperature from 37°C to 26°C overnight to destabilize  $\beta_2$  microglobulin and stripping the endogenous peptides from the cell using a mild acid treatment. The methods release previously bound peptides into the extracellular environment allowing new exogenous peptides to bind to the empty class I molecules. The cold-temperature incubation method enables exogenous peptides to bind efficiently to the MHC complex, but requires an overnight incubation at 26°C which may slow the cell's metabolic rate. It is also likely that cells not actively synthesizing MHC molecules

(e.g., resting PBMC) would not produce high amounts of empty surface MHC molecules by the cold temperature procedure.

Harsh acid stripping involves extraction of the peptides with trifluoroacetic acid, pH 2, or acid denaturation of the immunoaffinity purified class I-peptide complexes.

5 These methods are not feasible for CTL induction, since it is important to remove the endogenous peptides while preserving APC viability and an optimal metabolic state which is critical for antigen presentation. Mild acid solutions of pH 3 such as glycine or citrate-phosphate buffers have been used to identify endogenous peptides and to identify tumor associated T cell epitopes. The treatment is especially effective, in that only the  
10 MHC class I molecules are destabilized (and associated peptides released), while other surface antigens remain intact, including MHC class II molecules. Most importantly, treatment of cells with the mild acid solutions do not affect the cell's viability or metabolic state. The mild acid treatment is rapid since the stripping of the endogenous peptides occurs in two minutes at 4°C and the APC is ready to perform its function after  
15 the appropriate peptides are loaded. The technique is utilized herein to make peptide-specific APCs for the generation of primary antigen-specific CTL. The resulting APC are efficient in inducing peptide-specific CD8+ CTL.

Activated CD8+ cells may be effectively separated from the stimulator cells using one of a variety of known methods. For example, monoclonal antibodies  
20 specific for the stimulator cells, for the peptides loaded onto the stimulator cells, or for the CD8+ cells (or a segment thereof) may be utilized to bind their appropriate complementary ligand. Antibody-tagged molecules may then be extracted from the stimulator-effector cell admixture via appropriate means, e.g., via well-known immunoprecipitation or immunoassay methods.

25 Effective, cytotoxic amounts of the activated CD8+ cells can vary between *in vitro* and *in vivo* uses, as well as with the amount and type of cells that are the ultimate target of these killer cells. The amount will also vary depending on the condition of the patient and should be determined via consideration of all appropriate factors by the practitioner. Preferably, however, about  $1 \times 10^6$  to about  $1 \times 10^{12}$ , more preferably about  
30  $1 \times 10^8$  to about  $1 \times 10^{11}$ , and even more preferably, about  $1 \times 10^9$  to about  $1 \times 10^{10}$  activated CD8+ cells are utilized for adult humans, compared to about  $5 \times 10^6$  -  $5 \times 10^7$  cells used in mice.

Preferably, as discussed above, the activated CD8+ cells are harvested from the cell culture prior to administration of the CD8+ cells to the individual being treated. It is important to note, however, that unlike other present and proposed treatment modalities, the present method uses a cell culture system that is not tumorigenic.

5 Therefore, if complete separation of stimulator cells and activated CD8+ cells is not achieved, there is no inherent danger known to be associated with the administration of a small number of stimulator cells, whereas administration of mammalian tumor-promoting cells may be extremely hazardous.

10 Methods of re-introducing cellular components are known in the art and include procedures such as those exemplified in U.S. Patent No. 4,844,893 to Honsik, et al. and U.S. Patent No. 4,690,915 to Rosenberg. For example, administration of activated CD8+ cells via intravenous infusion is appropriate.

The immunogenic peptides of this invention may also be used to make monoclonal antibodies. Such antibodies may be useful as potential diagnostic or  
15 therapeutic agents.

The peptides may also find use as diagnostic reagents. For example, a peptide of the invention may be used to determine the susceptibility of a particular individual to a treatment regimen which employs the peptide or related peptides, and thus may be helpful in modifying an existing treatment protocol or in determining a prognosis  
20 for an affected individual. In addition, the peptides may also be used to predict which individuals will be at substantial risk for developing chronic infection.

To identify peptides of the invention, class I antigen isolation, and isolation and sequencing of naturally processed peptides was carried out as described in the related applications. These peptides were then used to define specific binding motifs  
25 for each of the following alleles A3.2, A1, A11, and A24.1. These motifs are described on page 3, above. The motifs described in Tables 8-11, below, are defined from pool sequencing data of naturally processed peptides as described in the related applications.

TABLE 8

## Summary

HLA-A3.2 Allele-Specific Motif (SEQ ID NO:378)

Position	Conserved Residues
1	-

2	V,L,M
3	Y,D
4	-
5	-
6	-
7	I
8	Q,N
9	K
10	K

TABLE 9

## Summary

HLA-A1 Allele-Specific Motif SEQ ID NO:218

Position	Conserved Residues
1	-
2	S,T
3	D,E
4	P
5	-
6	-
7	L
8	-
9	Y
10	K

TABLE 10

## Summary

HLA-A11 Allele-Specific Motif (SEQ ID NO:379)

Position	Conserved Residues
1	-
2	T,V

3	M,F
4	-
5	-
6	-
7	-
8	Q
9	K
10	K

TABLE 11

## Summary

HLA-A24.1 Allele-Specific Motif (SEQ ID NO:380)

Position	Conserved Residues
1	-
2	Y
3	I,M
4	D,E,G,K,P
5	L,M,N
6	V
7	N,V
8	A,E,K,Q,S
9	F,L
10	F,A

Example 25      Identification of immunogenic peptides

Using the motifs identified above for various MHC class I allele amino acid sequences from various pathogens and tumor-related proteins were analyzed for the presence of these motifs. Screening was carried out described in the related applications. Table 12 provides the results of searches of the antigens.

Table 12

Peptide	AA	Sequence	Source	A*0301	A*1101	SEQ ID NO:
28.0719	10	ILEQWVAGRK	HDV.nuc.16	0.0170	0.0012	219
28.0727	10	LSAGGKNLSK	HDV.nuc.115	0.0097	0.0150	220
1259.02	11	STDTVDTVLEK	Flu.HA.29	0.0001	0.0670	221
1259.04	9	GIAPLQLGK	Flu.HA.63	0.6100	0.2000	222
1259.06	10	VTAACSHAGK	Flu.HA.149	0.0380	0.0490	223
1259.08	9	GIHPSNSK	Flu.HA.195	0.1300	0.0140	224
1259.10	10	RMNYYWTLK	Flu.HA.243	2.5000	2.3000	225
1259.12	11	ITNKVNSVIEK	Flu.HA.392	0.0200	0.0670	226
1259.13	11	KMNIQFTAVGK	Flu.HA.402	0.0280	0.0092	227
1259.14	9	NIQFTAVGK	Flu.HA.404	0.0017	0.0330	228
1259.16	11	AVGKEFNKLEK	Flu.HA.409	0.0210	0.0460	229
1259.19	11	KVKSQLKNNAK	Flu.HA.465	0.0470	0.0031	230
1259.20	11	SVRNGTYDYPK	Flu.HA.495	0.0410	0.1400	231
1259.21	9	SIIPSGPLK	Flu.VMT1.13	0.7800	8.8000	232
1259.25	10	RMVLASTTAK	Flu.VMT1.178	0.5500	0.0350	233
1259.26	9	MVLASTTAK	Flu.VMT1.179	1.7000	1.4000	234
1259.28	10	RMGVQMQRFK	Flu.VMT1.243	0.1000	0.0059	235
1259.33	10	ATEIRASVGK	Flu.VNUC.22	0.1400	0.3000	236
1259.37	11	TMVMELVRMIK	Flu.VNUC.188	0.0890	0.0310	237
1259.43	10	RVLSFIKGTK	Flu.VNUC.342	0.8000	0.0830	238
F119.01	9	MSLQRQFLR	ORF3P	0.2000	0.7200	239
F119.02	9	LLGPGRPYR	TRP.197	0.0190	0.0091	240
F119.03	9	LLGPGRPYK	TRP.197K9	2.2000	0.6800	241
34.0019	8	RVYPELPK	CEA.139	0.0130	0.0440	242
34.0020	8	TVSAELPK	CEA.495	0.0037	0.0320	243
34.0021	8	TVYAEPPK	CEA.317	0.0160	0.0220	244
34.0029	8	TINYTLWR	MAGE2.74	0.0140	0.0550	245
34.0030	8	LVHFLLLK	MAGE2.116	0.0290	0.1500	246
34.0031	8	SVFAHPRK	MAGE2.237	0.1410	0.0810	247
34.0043	8	KVLHHMVK	MAGE3.285	0.0580	0.0190	248

34.0050	8	RVCACPGR	p53.273	0.3500	0.0490	249
34.0051	8	KMFCQLAK	p53.132	0.3800	0.3600	250
34.0062	8	RAHSSHLK	p53.363	0.5500	0.0071	251
34.0148	9	FVSNLATGR	CEA.656	0.0019	0.0490	252
34.0152	9	RLQLSNGNK	CEA.546	0.0250	0.0110	253
34.0153	9	RINGIPQQK	CEA.628	0.0400	0.0780	254
34.0154	9	KIRKYTMRK	HER2/neu.681	0.0620	0.0055	255
34.0155	9	LVHFLLLKK	MAGE2.116	0.5220	1.4000	256
34.0156	9	SMLEVFEGK	MAGE2.226	0.0950	1.6000	257
34.0157	9	SSFSTTINK	MAGE2.69	0.1600	2.0000	258
34.0158	9	TSYVKVLHK	MAGE2.281	0.5300	0.1500	259
34.0159	9	VIFSKASEK	MAGE2.149	0.4900	0.0530	260
34.0160	9	GSVVGWNQK	MAGE3.130	0.0040	0.2060	261
34.0161	9	SSLPTTMNK	MAGE3.69	0.6180	0.7100	262
34.0162	9	SVLEVFEGK	MAGE3.226	0.1330	0.9000	263
34.0171	9	SSBMGGMNK	p53.240	0.5440	1.1000	264
34.0172	9	SSCMGGMNK	p53.240	0.0090	0.0490	265
34.0211	10	RTLTLFNVTK	CEA.554	0.2200	1.3000	266
34.0212	10	TISPLNTSYK	CEA.241	0.1800	0.0330	267
34.0214	10	STTINYTLWK	MAGE2.72	0.0870	0.6500	268
34.0215	10	ASSLPTTMNK	MAGE3.68	0.0420	0.0270	269
34.0225	10	KTYQGSYGFK	p53.101	0.4900	0.4200	270
34.0226	10	VVRRBPHHEK	p53.172	0.1800	0.2100	271
34.0228	10	GLAPPQHLIK	p53.187	0.0570	0.0160	272
34.0229	10	NSSCMGGMNK	p53.239	0.0071	0.0290	273
34.0230	10	SSBMGGMNRK	p53.240	0.0420	0.1600	274
34.0232	10	RVCACPGRDK	p53.273	0.0190	0.0250	275
34.0295	11	KTITVSAELPK	CEA.492	0.3600	0.1600	276
34.0296	11	TTITVYAEPPK	CEA.314	0.0200	0.0280	277
34.0298	11	PTISPSYTYR	CEA.418	(0.0002)	0.1300	278
34.0301	11	GLLGDNQVMPK	MAGE2.188	0.0780	0.0047	279
34.0306	11	MVELVHFLLLK	MAGE2.113	0.0200	0.0120	280

34.0308	11	FSTTINYTLWR	MAGE2.71	0.0110	0.0170	281
34.0311	11	GLLGDNQIMPK	MAGE3.188	0.1300	0.0570	282
34.0317	11	RLGFLHSGTAK	p53.110	0.0430	0.0001	283
34.0318	11	ALNKMFCQLAK	p53.129	0.4400	0.0420	284
34.0323	11	RVCACPGRDRR	p53.273	0.0290	0.0290	285
34.0324	11	LSQETFSDLWK	p53.14	(0.0009)	0.0470	286
34.0328	11	RAHSSHLKSKK	p53.363	0.0270	0.0038	287
34.0329	11	VTCTYSPALNK	p53.122	0.0700	0.1200	288
34.0330	11	GTRVRAMAIYK	p53.154	1.1000	0.3300	289
34.0332	11	STSRHKKLMFK	p53.376	0.3100	0.1300	290
40.0107	9	LAARNVLVK	Her2/neu.846	0.0580	0.0285	291
40.0109	9	MALESILRR	Her2/neu.889	0.0034	0.0237	292
40.0145	10	ISWLGLRSLR	Her2/neu.450	0.0410	0.0027	293
40.0147	10	GSGAFGTVYK	Her2/neu.727	0.0660	0.1300	294
40.0153	10	ASPLDSTFYR	Her2/neu.997	0.0003	0.0670	295

### Example 3

#### Identification of immunogenic peptides

5                    Using the B7-like supermotifs identified in the related applications described above, sequences from various pathogens and tumor-related proteins were analyzed for the presence of these motifs. Screening was carried out described in the related applications. Table 13 provides the results of searches of the antigens.

10                    Table 13

Peptide	Sequence	Source	SEQ ID NO:
40.0013	SPGLSAGI	CEA.680I8	296
15 40.0022	KPYDGIPA	Her2/neu.921	297
40.0023	KPYDGIPI	Her2/neu.921I8	298
40.0050	APRMPEAA	p53.63	299
40.0051	APRMPEAI	p53.63I8	300



	40.0055	APAAPTPI	p53.76I8	301
	40.0057	APTPAAPI	p53.79I8	302
	40.0059	TPAAPAPI	p53.81I8	303
	40.0061	APAPAPSI	p53.84I8	304
5	40.0062	SPALNKMF	p53.127	305
	40.0063	SPALNKMI	p53.127I8	306
	40.0117	SPSAPPHRI	CEA.3I9	307
	40.0119	PPHRWCIPi	CEA.7I9	308
	40.0120	GPAYSGREI	CEA.92	309
10	40.0156	MPNQAQMRILI	Her2/neu.706I10	310
	40.0157	MPYGCLLDHVI	Her2/neu.801I10	311
	40.0161	APPHRWCIPIW	CEA.6	312
	40.0162	APPHRWCIPI	CEA.6I10	313
	40.0163	IPWQRLLTA	CEA.13	314
15	40.0164	IPWQRLLTI	CEA.13I10	315
	40.0166	LPQHILFGYSI	CEA.58I10	316
	40.0201	RPRFRELVSEF	Her2/neu.966	317
	40.0202	RPRFRELVSEI	Her2/neu.966I11	318
	40.0205	PPSPREGPLPA	Her2/neu.1149	319
20	40.0206	PPSPREGPLPI	Her2/neu.1149I11	320
	40.0207	GPLPAARPAGA	Her2/neu.1155	321
	40.0208	GPLPAARPAGI	Her2/neu.1155I11	322
	40.0231	APAPAAPTPAA	p53.74	323
	40.0232	APAPAAPTPAI	p53.74I11	324
25	40.0233	APAAPTPAAPA	p53.76	325
	40.0234	APAAPTPAAPI	p53.76I11	326
	45.0003	IPWQRLLI	CEA.13.I8	327
	45.0004	LPQHILFGI	CEA.58.I8	328
	45.0007	RPGVNLSI	CEA.428.I8	329
30	45.0010	IPQQHTQI	CEA.632.I8	330
	45.0011	TPNNNGTI	CEA.646.I8	331
	45.0016	CPLHNQEI	Her2/neu.315.I8	332
	45.0017	KPCARVCI	Her2/neu.336.I8	333

	45.0019	WPDSLPI	Her2/neu.415.I8	334
	45.0023	SPYVSRLI	Her2/neu.779.I8	335
	45.0024	VPIKWMAI	Her2/neu.884.I8	336
	45.0026	RPRFRELI	Her2/neu.966.I8	337
5	45.0028	APGAGGMI	Her2/neu.1036.I8	338
	45.0031	SPGKNGVI	Her2/neu.1174.I8	339
	45.0037	SPQGASSI	MAGE3.64.I8	340
	45.0038	YPLWSQSI	MAGE3.77.I8	341
	45.0044	SPLPSQAI	p53.33.I8	342
10	45.0046	MPEAAPPI	p53.66.I8	343
	45.0047	APAPSWPI	p53.86.I8	344
	45.0051	KPVEDKDAI	CEA.155.I9	345
	45.0054	IPQQHTQVI	CEA.632.I9	346
	45.0060	APPVAPAPI	p53.70.I9	347
15	45.0062	APAAPTPAI	p53.76.I9	348
	45.0064	PPGTRVRAI	p53.152.I9	349
	45.0065	APPQHLIRI	p53.189.I9	350
	45.0071	IPQQHTQVLI	CEA.632.I10	351
	45.0072	SPGLSAGATI	CEA.680.I10	352
20	45.0073	SPMCKGSRCI	Her2/neu.196.I10	353
	45.0074	MPNPEGRTYI	Her2/neu.282.I10	354
	45.0076	CPLHNQEVTI	Her2/neu.315.I10	355
	45.0079	KPDLSYMPII	Her2/neu.605.I10	356
	45.0080	TPSGAMPNQI	Her2/neu.701.I10	357
25	45.0084	GPASPLDSTI	Her2/neu.995.I10	358
	45.0091	APPVAPAPAI	p53.70.I10	359
	45.0092	APAPAAPTPI	p53.74.I10	360
	45.0093	APTPAAPAPI	p53.79.I10	361
	45.0094	APSWPLSSSI	p53.88.I10	362
30	45.0103	APTISPLNTSI	CEA.239.I11	363
	45.0108	SPSYTYRPGI	CEA.421.I11	364
	45.0117	CPSGVKPDLSI	Her2/neu.600.I11	365
	45.0118	SPLTSIISAVI	Her2/neu.649.I11	366

	45.0119	IPDGENVKIPI	Her2/neu.740.I11	367
	45.0124	SPLDSTFYRSI	Her2/neu.998.I11	368
	45.0128	LPAARPAGATI	Her2/neu.1157.I11	369
	45.0134	HPRKLLMQDLI	MAGE2.241.I11	370
5	45.0135	GPRALIETSYI	MAGE2.274.I11	371
	45.0139	GPRALVETSYI	MAGE3.274.I11	372
	45.0140	APRMPEAAPPI	p53.63.I11	373
	45.0141	VPSQKTYQGSI	p53.97.I11	374
	1145.10	FPHCLAFAY	HBV POL 541 analog	375
10	1145.09	FPVCLAFSY	HBV POL 541 analog	376
	26.0570	YPALMPLYACI	HBV.pol.645	377

The above description is provided to illustrate the invention but not to limit its scope. Other variants of the invention will be readily apparent to one of ordinary skill in the art and are encompassed by the appended claims. All publications, patents, and patent applications cited herein are hereby incorporated by reference.